

Antimicrobial Activity of Extracts of Bark of *Cinnamomum cassia* and *Cinnamomum zeylanicum*

Jasvin Singh¹, Rasvin Singh¹, Subramani Parasuraman^{2*}, Sathasivam Kathiresan¹

¹Department of Biotechnology, Faculty of Applied Sciences, AIMST University, Kedah, MALAYSIA.

²Unit of Pharmacology, Faculty of Pharmacy, AIMST University, Kedah, MALAYSIA.

ABSTRACT

Objective: To investigate the antimicrobial activities of various solvent extracts of the bark of *Cinnamomum cassia* and *Cinnamomum zeylanicum*.

Methods: The powdered barks of *C. cassia* and *C. zeylanicum* were extracted using methanol, ethanol and acetone. The antimicrobial activity of methanolic, ethanolic and acetone extracts of barks of *C. cassia* and *C. zeylanicum* were studied using agar well diffusion method against seven ATCC bacterial species (*Staphylococcus aureus*, *Bacillus cereus*, *Enterococcus faecalis*, *Streptococcus pyogenes*, *Escherichia coli*, *Salmonella bongori* and *Pseudomonas aeruginosa*). The antimicrobial activity of extracts of *C. cassia* and *C. zeylanicum* were compared with gentamycin. **Results:** The ethanolic and acetone extract of *C. cassia* and *C. zeylanicum* showed greater zones of inhibition than methanolic extract. *C. zeylanicum* inhibited the growth of all seven ATCC strains used in this study whereas, *C. cassia* inhibited the growth of *S. pyogenes*, *S. aureus*, *B. cereus*, *E. faecalis*, *P. aeruginosa* and *S. bongori*. Gentamycin inhibited

the pathogenic group of all ATCC strains used in this study. **Conclusion:** The ethanolic and acetone extracts of bark of *C. Cassia* and *C. zeylanicum* exhibited a broad spectrum of exhibited a broad-spectrum antimicrobial activity.

Key words: Agar well-diffusion method, Cinnamomum, Gentamycin, Minimum Inhibitory Concentration (MIC).

Correspondence

Dr. Subramani Parasuraman,

Unit of Pharmacology, Faculty of Pharmacy, AIMST University, Kedah, MALAYSIA.

Phone:+604-429 8108

Email: parasuphd@gmail.com

ORCID: <http://orcid.org/0000-0002-3729-5303>

DOI: 10.5530/ijpi.2020.2.26

INTRODUCTION

Antimicrobials are “any substance of natural, semisynthetic or synthetic origin that kills or inhibits the growth of microorganisms but causes little or no damage to the host”.¹ Antimicrobials are linked to adverse effects on consumption by the host which includes allergic reaction and hypersensitivity. In addition to this problem, commercialized antimicrobial drugs have caused multiple drug resistance in hosts and this is a major concern. Due to the evolution of bacteria being resistant towards modern medicine, this has forced to search for a new therapeutic agent that could be effective towards these bacteria. Therefore, it was a necessity to search an alternative drug derived from plants that could treat these diseases without triggering other side effects.² The benefits obtained from these medicinal plants are due to their secondary metabolites. These secondary metabolites are steroids, tannins, flavonoids, phenols, alkaloids and these compounds are present in different quantities depending on the part of the plant.^{3,4}

Cinnamomum cassia and *Cinnamomum zeylanicum* (family: Lauraceae) are the well-known plant species for its medicinal properties. *C. cassia* also called Chinese cassia is widely cultivated in China. The dried stem bark of *C. cassia* is one of the important food spices and also has considered having medicinal properties, such as antimicrobial, anti-tumorigenic, anti-inflammatory and antidiabetic characteristics.⁵ *C. cassia* is commonly used as traditional Chinese medicine for treating dyspepsia, gastritis, blood circulation disturbances and inflammatory diseases.⁶ *C. zeylanicum* also called Ceylon cinnamon is widely cultivated in Sri Lanka and has antidiabetic, anti-inflammatory, antibacterial, antiviral and antifungal activities.^{7,8} Cinnamon Oil and cinnamaldehyde from the *C. cassia* is showed antimicrobial activity against both gram-positive bacteria (*Staphylococcus aureus*) and gram-negative bacteria including *Escherichia coli*, *Enterobacter aerogenes*, *Proteus vulgaris*,

Pseudomonas aeruginosa, *Vibrio cholerae*, *Vibrio parahaemolyticus* and *Samonella typhymurium*.⁹ The essential oil from the bark of *C. zeylanicum* has antimicrobial activity and highly active against both gram-positive (*Staphylococcus*, *Streptococcus*, *Enterococcus*) and gram-negative bacteria including *Pseudomonas aeruginosa*.¹⁰ The antimicrobial activity of essential oil from *C. cassia* and *C. zeylanicum* is well documented. The antimicrobial activity of extracts of bark of *C. cassia* and *C. zeylanicum* is not clear. Hence the present study is planned to investigate the antimicrobial activities of various solvent extracts of bark of *C. cassia* and *C. zeylanicum*.

MATERIALS AND METHODS

Collection and identification of barks of *Cinnamomum*

The cinnamon barks were obtained from Penang Spice Garden, Pulau Pinang and the authenticity of the *C. cassia* and *C. zeylanicum* species was confirmed by Dr. Deivanai Subramanian. The bark was cleaned thoroughly and placed in an oven to be dried for 24 h at 40°C. The dried bark was then powdered using a blender and the powder was stored in an airtight container.

Extraction of *Cinnamomum* bark

The powdered barks of *C. cassia* and *C. zeylanicum* were packed in a thimble and placed in a Soxhlet apparatus and extracted using methanol, ethanol and acetone. The extraction was carried out for 24 h at about 55°C–80°C; the extract was filtered through a muslin cloth. The filtrate was concentrated to a dry mass by evaporation under reduced pressure. The extracts of *C. cassia* and *C. zeylanicum* were stored in a desiccator at room temperature until further analysis. The yield of methanolic,

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

ethanolic and acetone extract of barks of *C. cassia* was found to be ≈ 22 , ≈ 17 and $\approx 15\%$ w/v, respectively. The percentage yield of methanolic, ethanolic and acetone extract of *C. zeylanicum* was found to be ≈ 21 , ≈ 19 , and $\approx 12\%$ respectively.

Test organisms

Total of seven pure ATCC bacterial strains were selected and obtained from Bio-focus Saintifik Sdn Bhd with 4 strains being gram-positive (*Staphylococcus aureus*, *Bacillus cereus*, *Enterococcus faecalis*, and *Streptococcus pyogenes*) and the remaining 3 strains being gram-negative (*Escherichia coli*, *Salmonella bongori* and *Pseudomonas aeruginosa*).

Antimicrobial Activity of extracts of *C. cassia* and *C. zeylanicum*

Agar well-diffusion method used to evaluate the antimicrobial activity of an extract and this method is almost similar to the disk-diffusion method.¹¹ The surface of the Petri-plate containing agar is inoculated by streaking the specific bacterial strain over it with even distribution. Sterile cork-borer of the desired diameter is used to create wells on the agar in order for the extracts to be filled into these wells.

Bacterial strains were prepared using nutrient broth and kept in an incubator at 37°C for 24 hrs. Bacterial growth is observed in the nutrient broth which was then diluted to contain 10⁵ viable cells in accordance with the McFarland standards. A 25 g of Mueller-Hinton Agar (MHA) was dissolved into 200 ml of distilled water and was autoclaved at 121°C for 15 min at 15 Psi. The MHA was then poured into Petri-plates with a volume of 15 ml and left to cool. 150 μ l of the bacterial culture were spread equally on the plate and left to dry. 8 mm sterile cork-borer was used to create wells on the agar in order for the extracts ranging from 20–100 μ l to be filled into these wells. Four wells were created to fill 3 concentrations (0.5 g, 1 g and 1.5 g) as well as a well for the standard antibiotic gentamycin (1 mg). This process was repeated for the remaining 6 ATCC bacterial strains and the Petri-plates were properly labeled and stored in an incubator at 37°C for 24 hrs.¹² All tests were carried out in triplicates and the mean zones of inhibition were determined. The Minimum Inhibitory Concentration (MIC) was further determined.

Determination of Minimum Inhibitory Concentration

MIC of the extracts was carried out using the broth dilution method. MIC is said to be the lowest concentration of the sample, which inhibits the visible growth of microbes.¹³ Serial dilution was carried out accordingly with concentrations of 1000 mg/ml, 500 mg/ml, 250mg/ml, 125 mg/ml and 62.5 mg/ml. Mueller-Hinton (MH) broth was prepared and 1 ml was added into test tubes alongside 100 μ l of selected ATCC strains. Then, 1 ml of extracts with the concentrations mentioned above were introduced into the test tubes contained the broth and ATCC strain and mixed for even distribution. These test tubes were then incubated for 36 hrs at 37°C. As a positive control, 1 ml of selected antibiotics was added to the test tube while the negative control only contained selected ATCC strains. The results obtained after incubation were compared to both the positive and negative control. The concentration of the extract that gave clear results (absence of turbidity) was considered as the MIC of the extract.¹⁴

RESULTS

Preliminary antimicrobial activity of extracts of *C. cassia* and *C. zeylanicum*

Preliminary antimicrobial screening of extract of *C. cassia* (150 mg/ml) and *C. zeylanicum* (150 mg/ml) tested against *E. coli*, *P. aeruginosa*,

Table 1: Results of a preliminary study of the extracts against ATCC cultures.

ATCC Strains	<i>C. cassia</i>			<i>C. zeylanicum</i>		
	Methanol	Ethanol	Acetone	Methanol	Ethanol	Acetone
Gram positive bacteria						
<i>Streptococcus pyogenes</i>	/	/	/	/	/	/
<i>Staphylococcus aureus</i>	/	/	/	X	/	/
<i>Bacillus cereus</i>	/	/	/	/	X	/
<i>Enterococcus faecalis</i>	/	/	/	X	/	/
Gram negative bacteria						
<i>Escherichia coli</i>	X	X	X	X	/	/
<i>Pseudomonas aeruginosa</i>	/	/	/	/	/	/
<i>Salmonella bongori</i>	/	/	/	/	X	/

(/) indicates inhibitory action; (X) indicates absence of inhibitory action.

S. bongori, *S. pyogenes*, *S. aureus*, *B. cereus* and *E. faecalis*. The results of preliminary antimicrobial activity of extract of *C. cassia* and *C. zeylanicum* were summarized in Table 1.

Antimicrobial activity of extracts of *C. cassia* and *C. zeylanicum*

Following the preliminary screening, all the 7 ATCC strains were tested (in triplicates) using the agar-well method to determine its activity against different concentrations of the extract of *C. cassia* and *C. zeylanicum*. The antimicrobial activity of extracts of *C. cassia* and *C. zeylanicum* on gram-positive and gram-negative organism were presented in Table 2 and 3 respectively. The antimicrobial activity of extract of *C. cassia* and *C. zeylanicum* were compared with gentamycin. Overall, gentamycin inhibited the pathogenic group of all ATCC strains used in this study. The ethanolic and acetone extract of *C. cassia* and *C. zeylanicum* showed greater zones of inhibition than methanolic extract. *C. zeylanicum* inhibited the growth of all seven ATCC strains used in this study whereas, *C. cassia* inhibited the growth of *S. pyogenes*, *S. aureus*, *B. cereus*, *E. faecalis*, *P. aeruginosa* and *S. bongori*.

Minimum Inhibitory Concentration

MIC values (μ g/ml) of the different solvent extracts of *C. cassia* and *C. zeylanicum* against the ATCC bacterial strains were summarized in Table 4. The methanolic and ethanolic extracts of *C. cassia* showed turbidity in the broth dilution method whereas, acetone extract of *C. cassia* inhibited the bacterial growth (absence of turbidity) of few ATCC strains. The methanolic, ethanolic and acetone extracts of *C. zeylanicum* inhibited bacterial growth of few ATCC strains.

DISCUSSION

The bacterial strains used were of different taxonomy. In this study, both *C. cassia* and *C. zeylanicum* bark extracts showed antimicrobial activity against gram-positive and gram-negative bacteria. The antibacterial activity has been associated with the presence of active phytoconstituents such as phenols and flavonoids in the bark extract.¹⁵ Cinnamon (*C. zeylanicum* and *C. cassia*) primarily contains vital oils and other derivatives, such as cinnamaldehyde, cinnamic acid and cinnamate. Cinnamaldehyde and trans-cinnamaldehyde are the most important constituents of cinnamon.¹⁶ The antibacterial activity of cinnamon is said to be associated with its major component, cinnamaldehyde. This compound inhibits bacterial acetyl-CoA carboxylase.¹⁷ Acetyl-CoA carboxylase (ACC) functions by catalyzing the first important step in

Table 2: Antimicrobial activity of extracts of *C. cassia* and *C. zeylanicum* against gram-positive bacteria.

Test compound	concentration	Zone of Inhibition (mm)			
		<i>S. pyogenes</i>	<i>S. aureus</i>	<i>B. cereus</i>	<i>E. faecalis</i>
Gentamycin	1mg/ml	26.75 ± 0.57	16.94 ± 0.39	21.72 ± 0.58	22.16 ± 0.35
Methanolic extract of <i>C. cassia</i>	50 mg/ml	-	-	-	-
	100 mg/ml	21.5 ± 1.32	-	13.8 ± 1.26	16.0 ± 0.87
	150 mg/ml	21.7 ± 2.08	18.5 ± 1.32	17.7 ± 1.26	18.3 ± 1.53
Ethanol extract of <i>C. cassia</i>	50 mg/ml	-	-	-	-
	100 mg/ml	17.8 ± 1.04	11.7 ± 1.26	16.7 ± 1.26	16.5 ± 1.50
	150 mg/ml	21.8 ± 1.61	16.5 ± 1.32	18.5 ± 0.50	18.2 ± 0.29
Acetone extract of <i>C. cassia</i>	50 mg/ml	-	9.8 ± 1.04	-	-
	100 mg/ml	24.0 ± 1.0	22.0 ± 1.00	22.0 ± 2.00	16.2 ± 1.04
	150 mg/ml	28.0 ± 0.87	23.7 ± 1.26	26.2 ± 1.04	24.7 ± 0.29
Methanolic extract of <i>C. zeylanicum</i>	50 mg/ml	-	-	-	-
	100 mg/ml	13.16 ± 0.46	-	13.26 ± 0.59	-
	150 mg/ml	16.32 ± 0.71	-	14.21 ± 0.73	-
Ethanol extract of <i>C. zeylanicum</i>	50 mg/ml	-	18.53 ± 0.96	-	-
	100 mg/ml	15.15 ± 1.00	19.54 ± 0.58	-	-
	150 mg/ml	16.34 ± 0.80	20.48 ± 0.53	-	13.51 ± 0.56
Acetone extract of <i>C. zeylanicum</i>	50 mg/ml	-	-	15.65 ± 0.34	-
	100 mg/ml	17.41 ± 0.94	-	15.70 ± 0.57	10.10 ± 0.46
	150 mg/ml	20.48 ± 0.51	17.31 ± 0.33	16.31 ± 0.70	14.34 ± 0.65

All the values are Mean ± SD (n = 3).

Table 3: Antimicrobial activity of extracts of *C. cassia* and *C. zeylanicum* against gram-negative bacteria.

Test compound	concentration	Zone of Inhibition (mm)		
		<i>E. coli</i>	<i>P. aeruginosa</i>	<i>S. bongori</i>
Gentamycin	1mg/ml	16.64 ± 0.54	29.27 ± 0.67	20.72 ± 1.51
Methanolic extract of <i>C. cassia</i>	50 mg/ml	-	-	-
	100 mg/ml	-	22.7 ± 0.76	16.1 ± 0.76
	150 mg/ml	-	23.5 ± 1.32	20.5 ± 1.80
Ethanol extract of <i>C. cassia</i>	50 mg/ml	-	-	-
	100 mg/ml	-	28.2 ± 1.04	16.7 ± 1.26
	150 mg/ml	-	28.7 ± 1.53	19.7 ± 2.08
Acetone extract of <i>C. cassia</i>	50 mg/ml	-	16.0 ± 1.73	21.8 ± 1.04
	100 mg/ml	-	27.8 ± 1.61	24.8 ± 1.04
	150 mg/ml	-	32.3 ± 2.08	26.0 ± 1.00
Methanolic extract of <i>C. zeylanicum</i>	50 mg/ml	-	-	11.22 ± 0.85
	100 mg/ml	-	-	14.23 ± 0.43
	150 mg/ml	-	14.27 ± 1.05	16.14 ± 0.80
Ethanol extract of <i>C. zeylanicum</i>	50 mg/ml	17.31 ± 1.13	-	-
	100 mg/ml	21.15 ± 1.03	14.40 ± 0.67	-
	150 mg/ml	22.56 ± 0.96	17.14 ± 1.01	-
Acetone extract of <i>C. zeylanicum</i>	50 mg/ml	-	-	11.12 ± 0.87
	100 mg/ml	10.91 ± 0.74	-	11.33 ± 1.28
	150 mg/ml	12.74 ± 0.58	16.20 ± 1.05	17.77 ± 0.75

All the values are Mean ± SD (n = 3).

Table 4: Minimum Inhibitory Concentration results of the plant extract (µg/ml).

ATCC bacterial strains	<i>C. cassia</i>			<i>C. zeylanicum</i>		
	Methanol	Ethanol	Acetone	Methanol	Ethanol	Acetone
<i>E. coli</i>	-	-	-	-	250	125
<i>S. bongori</i>	-	-	500	125	-	125
<i>S. pyogenes</i>	-	-	-	125	125	250
<i>S. aureus</i>	-	-	500	-	250	-
<i>P. aeruginosa</i>	-	-	500	-	125	-
<i>B. cereus</i>	-	-	-	125	-	250
<i>E. faecalis</i>	-	-	-	-	-	125

fatty acid biosynthesis which is a metabolic pathway required for several important biological processes which include the synthesis and maintenance of cellular membranes.¹⁸

The antimicrobial activity of these extracts is said to be to arise principally from the potential hydrophobicity of the extracts in obstructing the proper formation of the bacterial cell membrane and its other important structures which in return leads to ion leakage. It has been suggested that eugenol and cinnamaldehyde inhibit the production of an important and essential enzyme produced by the bacteria which in return causes damage to the cell wall of the bacteria.¹⁹ This could be explained by the principle of their hydrophobicity which may allow the separation of the lipids of the bacterial cell membrane which in return would make them more permeable hence leading to leakage of ions and other cell constituents.^{20,21} The bark of cinnamon contains 65 to 80% of cinnamaldehyde and 5 to 10% of eugenol.¹⁶

The different parts of cinnamon plants studied for their antimicrobial activity. Matan *et al.* reported the effects of cinnamon oils on different bacterial (*Pediococcus halophilus* and *Staphylococcus aureus*), fungal (*Aspergillus flavus*, *Mucor plumbeus*, *Penicillium roqueforti* and *Eurotium* sp.) and yeast species (*Candida lipolytica*, *Pichia membranaefaciens*, *Debaryomyces hansenii* and *Zygosaccharomyces rouxii*).²² Varalakshmi *et al.* reported the effects of bark extract of *C. zeylanicum* on different bacterial strains such as *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*. The findings of this study showed that the bark extracts of *C. zeylanicum* inhibiting both gram-positive bacteria and gram-negative bacteria.²³ These study result indicating that cinnamon is a natural antimicrobial agent with a broad spectrum of activity.

The major limitation of the study is that the method of antimicrobial assay. The method of antimicrobial study may affect the result.²⁴ Also, factors such as choice of bacterial strains selected for the study, their sensitivity, incubation time, as well as temperature which affect the result.^{18,24}

CONCLUSION

In conclusion, the results of this investigation revealed the ethanolic and acetone extract of *C. cassia* and *C. zeylanicum* showed greater zones of inhibition than methanol extract. *C. zeylanicum* inhibited the growth of all seven ATCC strains used in this study whereas, *C. cassia* inhibited the growth of *S. pyogenes*, *S. aureus*, *B. cereus*, *E. faecalis*, *P. aeruginosa* and *S. bongori*. The bark of *C. Cassia* and *C. zeylanicum* exhibited a broad-spectrum antimicrobial activity.

ACKNOWLEDGEMENT

Nil.

CONFLICT OF INTEREST

The author declares no conflict of interest.

ABBREVIATIONS

ACC: Acetyl-CoA carboxylase; MH: Mueller- Hinton; MHA: Mueller-Hinton Agar; MIC: Minimum Inhibitory Concentration.

REFERENCES

- Mariottini GL, Grice ID. Antimicrobials from Cnidarians. A New Perspective for Anti-Infective Therapy?. Mar Drugs. 2016;14(3):E48.
- Agrawal P, Rai V, Singh RB. Randomized placebo-controlled, single blind trial of holy basil leaves in patients with noninsulin-dependent diabetes mellitus. Int J Clin Pharmacol Ther. 1996;34(9):406-9.
- Joshi AR, Edington JM. The use of medicinal plants by two village communities in the central development region of Nepal. Econ Bot. 1990;44(1):71-83.
- Manandhar NP. Traditional medicinal plants used by tribals of Lamjung District, Nepal. Int J Crude Drug Res. 1987;25(4):236-40.
- Chou ST, Chang WL, Chang CT, Hsu SL, Lin YC, Shih Y. *Cinnamomum cassia* essential oil inhibits α -MSH-induced melanin production and oxidative stress in murine B16 melanoma cells. Int J Mol Sci. 2013;14(9):19186-201.
- Liao JC, Deng JS, Chiu CS, Hou WC, Huang SS, Shie PH, *et al.* Anti-inflammatory activities of *Cinnamomum cassia* constituents *in vitro* and *in vivo*. Evid Based Integr Med Complement Altern Med. 2012.
- Paranagama PA, Wimalasena S, Jayatilake GS, Jayawardena AL, Senanayake UM, Mubarak AM. A comparison of essential oil constituents of bark, leaf, root and fruit of cinnamon (*Cinnamomum zeylanicum* Blum) grown in Sri Lanka. J Natl Sci Found Sri. 2001;29(3-4):147-53.
- Simic A, Sokovic MD, Ristic M, Grujic-Jovanovic S, Vukojevic J, Marin PD. The chemical composition of some Lauraceae essential oils and their antifungal activities. Phytother Res. 2004;18(9):713-7.
- Ooi LS, Li Y, Kam SL, Wang H, Wong EY, Ooi VE. Antimicrobial activities of cinnamon oil and cinnamaldehyde from the Chinese medicinal herb *Cinnamomum cassia* Blume. Am J Chin Med. 2006;34(03):511-22.
- Unlu M, Ergene E, Unlu GV, Zeytinoglu HS, Vural N. Composition, antimicrobial activity and *in vitro* cytotoxicity of essential oil from *Cinnamomum zeylanicum* Blume (Lauraceae). Food Chem Toxicol. 2010;48(11):3274-80.
- Drago L, DeVecchi E, Nicola L, Gismondo MR. *In vitro* evaluation of antibiotics' combinations for empirical therapy of suspected methicillin resistant *Staphylococcus aureus* severe respiratory infections. BMC Infect Dis. 2007;7(1):111.
- Okudoh VI, Wallis FM. Antimicrobial activity of rare actinomycetes isolated from natural habitats in KwaZulu-Natal, South Africa. South Afr J Sci. 2007;103(5-6):216-22.
- Furletti VF, Teixeira IP, Obando-Pereda G, Mardegan RC, Sartoratto A, Figueira GM, *et al.* Action of *Coriandrum sativum* L. essential oil upon oral *Candida albicans* biofilm formation. Evid Based Complement Altern Med. 2011;2011:985832.
- Yu T, Lee S, Yang WS, Jang HJ, Lee YJ, Kim TW, *et al.* The ability of an ethanol extract of *Cinnamomum cassia* to inhibit Src and spleen tyrosine kinase activity contributes to its anti-inflammatory action. J Ethnopharmacol. 2012;139(2):566-73.
- Yamamoto S, Gaynor C. Therapeutic potential of inhibitory of the NF- κ B pathway in the treatment of inflammation and cancer. J Clin Investig. 2002;1:493-503.
- Rao PV, Gan SH. Cinnamon: A multifaceted medicinal plant. Evid Based Complement Alternat Med. 2014;642942.

17. Meades G, Henken RL, Waldrop GL, Rahman MM, Gilman SD, Kamatou GP, Viljoen AM, et al. Constituents of cinnamon inhibit bacterial acetyl CoA carboxylase. *Planta Med.* 2010;76(14):1570-5.
18. Gago G, Diacovich L, Arabolaza A, Tsai SC, Gramajo H. Fatty acid biosynthesis in actinomycetes. *FEMS Microbiol Rev.* 2011;35(3):475-97.
19. Burt S. Essential oils: their antibacterial properties and potential applications in foods-a review. *International Journal of Food Microbiology.* 2004;94(3):223-53.
20. Teke GN, Elisée KN, Roger KJ. Chemical composition, antimicrobial properties and toxicity evaluation of the essential oil of *Cupressus lusitanica* Mill. leaves from Cameroon. *BMC Complement Altern Med.* 2013;13(1):130.
21. Ultee A, Bennik MH, Moezelaar RJ. The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Appl Environ Microbiol.* 2002;68(4):1561-8.
22. Matan N, Rimkeeree H, Mawson AJ, Chompreeda P, Haruthaitanasan V, Parker M. Antimicrobial activity of cinnamon and clove oils under modified atmosphere conditions. *Int J Food Microbiol.* 2006;107(2):180-5.
23. Varalakshmi B, Anand AV, Karpagam T, Bai JS, Manikandan R. *In vitro* antimicrobial and anticancer activity of *Cinnamomum zeylanicum* Linn bark extracts. *Int J Pharm Pharm Sci.* 2014;6(1):12-8.
24. Mith H, Dure R, Delcenserie V, Zhiri A, Daube G, Clinquart A. Antimicrobial activities of commercial essential oils and their components against food-borne pathogens and food spoilage bacteria. *Food Sci Nutr.* 2014;2(4):403-16.
25. Bozin B, Mimica-Dukic N, Simin N, Anackov G. Characterization of the volatile composition of essential oils of some Lamiaceae spices and the antimicrobial and antioxidant activities of the entire oils. *J Agric Food Chem.* 2006;54(5):1822-8.

Article History: Submission Date : 28-03-2020; Revised Date : 15-04-2020; Acceptance Date : 02-05-2020.

Cite this article: Singh J, Singh R, Parasuraman S, Kathiresan S. Antimicrobial Activity of Extracts of Bark of *Cinnamomum Cassia* and *Cinnamomum zeylanicum*. *Int. J. Pharm. Investigation.* 2020;10(2):141-5.