

Altitude Variation in Volatile Composition of Blueberry Leaf Analyzed by SPME GC-MS

İshak Erik¹, Gözde Kılıç¹, Sevde Seher Yaylı², Gülşah Yılmaz², Kamil Coşkunçelebi², Nurettin Yaylı^{1,*}

¹Department of Pharmacognosy, Faculty of Pharmacy, Karadeniz Technical University, Trabzon, TURKEY.

²Department of Biology, Faculty of Science, Karadeniz Technical University, Trabzon, TURKEY.

ABSTRACT

Background: The present work was aimed to carry out volatile component analyzes in *Vaccinium arctostaphylos* L., *V. uliginosum* L., *V. vitis-idaea* L. and *V. myrtillus* L leaf growing at different altitudes of the East Blacksea Region of Turkey. **Methods:** The leaf of *Vaccinium* species were harvested from twenty-one different altitudes (748-3035 m) from six cities (Artvin-Ardahan-Rize-Trabzon-Gümüşhane-Giresun) of Turkey. The diversity of volatiles in the leaf was investigated by SPME GC-FID/MS. **Results:** The major constituents of the *Vaccinium* leaf showed variation with changes in altitudes. The identified volatile components of *V. arctostaphylos* and *V. uliginosum* were represented mainly by aldehydes in all altitudes, whereas monoterpenes were found the major constituent of *V. vitis-idaea* at Posof-Ardahan (2376 m) and Artvin (2553 m) samples. In all altitudes, capronaldehyde (7.23-28.96%) and 2(E)-hexenal (8.90-53.59%) in the leaf of *V. arctostaphylos*; capronaldehyde (17.04-37.09%) and limonene (16.50-47.51%) in the leaf of *V. vitis-idaea* and capronaldehyde (4.55-39.90%), 2(E)-hexenal (25.08-80.99%) and hexadecane (2.97-11.32%) in the leaf of *V. uliginosum*; and capronaldehyde (14.66-37.26%) and 2(E)-hexenal

(18.18-37.59%) at the altitudes of 1912m, 2533 m and 2565 m in the leaf of *V. myrtillus* were the major constituents with the different percentages, respectively. 2-Bornanone (32.86%) at the altitude of 2613 m and 3-penten-2-one (93.60%) at the altitude of 2811m in the leaf of *V. myrtillus* were found to be the major compounds. **Conclusion:** Comparisons of the volatile components of *V. arctostaphylos*, *V. uliginosum*, *V. vitis-idaea* and *V. myrtillus* in different sites showed significant differences among populations at different altitudes.

Key words: *Vaccinium arctostaphylos*, *V. uliginosum*, *V. vitis-idaea*, *V. myrtillus*, SPME GC-FID/MS.

Correspondence

Prof. Nurettin Yaylı

Department of Pharmacognosy, Faculty of Pharmacy, Karadeniz Technical University, Trabzon, TURKEY.

Phone no: +90-05333153187

Email: yayli@ktu.edu.tr

DOI: 10.5530/ijpi.2020.4.90

INTRODUCTION

The genus *Vaccinium* L. (Ericaceae) is represented approximately with 450 species of shrubs or small trees worldwide.¹ Four of the species (*V. arctostaphylos* (Va), *V. uliginosum* (Vu), *V. vitis-idaea* (Vvi) and *V. myrtillus* (Vm)) occur in Turkey and these are mostly distributed in North-East and North-West Anatolia.² *Vaccinium* is a medicinal plant that leaf consist on deciduous or evergreen and berries are vary from blue, purple, black or red. Berries can be eaten fresh; leaf can be dried and steeped. Also *Vaccinium* leaf, flower, berries and rhizomes are more usually processed.³ The genus is rich sources of dietary anthocyanins and antioxidants. Leaf and berries are traditionally used in for the treatment of diabetes^{4,5} and also including antimicrobial, antiinflammatory and antimutagenic properties.^{6,7} *Vaccinium* species are an industrial medicinal plant with various pharmaceutical and nutritional applications. Major constituents of the essential oil of *Vaccinium* species were mentioned to be as; α -terpineol (14.99%) and linalool (13.7%) for Va,⁸ nerolidol (20.01%), (Z,Z,Z)-1,5,9,9-tetramethyl-1,4,7-cycloundecatriene (17.99%) and caryophyllene (9.59%) for *Vaccinium bracteatum* Thumb.⁹ α -terpineol (4.7%) and linalool (17.0%) for Vvi,¹⁰ α -pinene (15.5%), linalool (11.7%), sandaracopimaradiene (5.9%) and safranal (8.8%) for *V. arctostaphylos*;¹¹ diisobutyl phthalate for Vvi (17.96%);¹² decahydro dimethylnaphthalene (47.04%), linalool (3.19%), 3,7-dimethyl-1,5,7-octatrien-3-ol (1.40%) and benzene acetaldehyde (1.03%) for *Vaccinium dunalianum* var. *dunalianum* (C. B. Clarke) Ridley;¹³ α -pinene for *Vaccinium angustifolium* Aiton, (23%) and *Vaccinium arboreum* Marsh., (63%), methyl benzoate for *Vaccinium varingiaefolium* Miq., (18%) and

ethyl benzene for *Vaccinium poasanum* Donn. Sm. (21%);¹⁴ and butyl butanoate, *cis*-3-hexen-1-ol, α -terpineol, geraniol and *trans*-2-hexenal for *Vaccinium corymbosum* L.¹⁵ HS SPME GC-MS analysis for fruit juice of Vm were mentioned.¹⁶ The complexity of blueberry (Vm) aroma was explored by SPME GC-MS and VOCs were reported as aldehydes, alcohols, terpenoids and esters.¹⁷ Volatile compounds, the antifungal activity and the phytotoxic activity of essential oil of Vm were also reported.^{18,19} In the literature, polar metabolites of leaf of Vm and Vvi were also studied by HPLC-DAD, HPLC-MS and GC-FID,²⁰ and the extracts of entire fruits and leaf of Vm collected in Finland and Poland were quantitated by GC-FID/MS and the main bilberry constituents were given as α - and β -amyrin, α - and β -amyrenone, campesterol, cholesterol and citrostadienol.²¹ However, to the best of our knowledge of literature survey, no data about altitude variation in volatile composition of Va, Vu, Vvi and Vm leaf analyzed by SPME GC-MS have been reported up to date. The present study aim to test whether there is any altitude effect on volatile compounds of Va, Vu, Vvi and Vm leaf growing spontaneously in different altitudes in the East Blacksea region of Turkey.

MATERIALS AND METHODS

Plant materials

Va, Vu, Vvi and Vm leaf (25 g, wet) were harvested from different altitudes in the East Blacksea Region (Artvin-Ardahan-Rize-Trabzon-Gümüşhane-Giresun) of Turkey (Table 1) and were identified.^{2,3}

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Vouchers were deposited in the Herbarium of Biology (KTUB) at Karadeniz Technical University, Turkey (Table 1).

Solid Phase Micro Extraction (SPME)

The blended leaf (2 g) were placed to a sealed SPME vial (10 mL) with a silicone-rubber septum cap then submitted to solid-phase micro extraction device (Supelco, USA). A DVB/Carboxen/PDMS coating fiber was used to obtain volatile components. The SPME fibers were conditioned for 5 min at 250°C in the GC injector. Extraction was achieved with magnetic stirring at 80°C using an incubation time of 5 min and an extraction time of 10 min. Fiber with extract of volatile compounds were subsequently injected into the GC injector. Each sample was analyzed and reported. Conditioning time for subsequent assays was set at 4 min of desorption after each extraction. The temperature, incubation and extraction times were set according to the reported experiment.²²⁻²⁴

Gas chromatography-Mass spectrometry (GC-FID/MS)

The gas chromatography-flame ionization detector (GC-FID) analysis was carried out on a Shimadzu QP2010 plus gas chromatography equipped with a flame ionization detector (FID) using a Rtx-5MS capillary column (30 m x 0.25 mm, film thickness, 0.25 µm). Shimadzu QP2010 Plus gas chromatograph was coupled to a Shimadzu QP2010 Ultra mass selective detector. The injection was performed in split mode

(1:30) at 230°C. The column held initially at 60°C for 2 min and then increased to 240°C with a 3°C/min heating ramp. The oven program was as follows: the initial temperature was 60°C for 2 min, which was increased to 240°C at 3 min, the final temperature of 250°C was held for 4 min. Helium (99.999 %) was used as carrier gas with a constant flow-rate of 1 mL/min. Detection was implemented in electronic impact mode (EI); ionization voltage was fixed at 70 eV, scan mode (40-450 *m/z*) was used for mass acquisition.²²⁻²⁵ Each sample was analyzed and mean reported.

Identification of Volatile Constituents

Retention indices of the volatile components of leaf belong to four *Vaccinium* species were determined by Kovats method using *n*-alkanes (C₆-C₃₂) as standards.²⁶ Volatile compounds were identified by comparing mass spectra and Kovats' Retention Indices with those in literature²²⁻³¹ and by computer searching followed by matching the mass spectra data with those held in computer library (NIST, Wiley7NL, FFNSC1.2 and W9N11).

RESULTS

The volatile compounds of leaf belong to four *Vaccinium* species collected from different altitudes in the East Blacksea Region of Turkey were analyzed by SPME GC-FID/MS. A total of 16, 23, 13, 21, 16,

Table 1: Examined species, voucher numbers, collection data and localities of the *Vaccinium* species in Turkey

Sample	Harvested localities	Altitude (m)	Harvested date	Collection number ^a
<i>Vaccinium arctostaphylos</i> (Va)				
Va1	A8 Rize: Ardeşen, Işıklı village, path route, forestry region	748	19 viii 2017	C&Y-46
Va2	A8 Rize: Ardeşen, Siprona plateau, forestry regions	1032	19 viii 2017	C&Y-47
Va3	A7 Trabzon: Araklı, Köyiçi village, forestry regions	1443	06 vii 2017	C&Y-15
Va4	A7 Trabzon: Düzköy, Çayırbağı, Kale river, forestry regions	1516	06 vii 2017	C&Y-21
Va5	A7 Giresun: Doğankent, Harmançık plateau, forestry regions	1666	02 vii 2017	C&Y-14
Va6	A7 Gümüşhane: Kürtün above plateau, forestry regions	1845	01 vii 2017	C&Y-1
Va7	A8 Artvin: Şavşat, Pınarlı plateau, forestry regions	2105	07 ix 2018	C&Y-117
Va8	A8 Artvin: Murgul, Tiryol mountain, forestry regions	2234	10 vii 2018	C&Y-91
<i>Vaccinium uliginosum</i> (Vu)				
Vu1	A8 Artvin: Şavşat, Karagöl National Park, alpine meadows	2530	02 vii 2018	C&Y-95
Vu2	A8 Rize: İkizdere, Anzer plateau, alpine meadows	2537	02 ix 2018	C&Y-108
Vu3	A9 Ardahan: Hanak, Alabalık plateau, alpine meadows	2588	08 vii 2018	C&Y-101
Vu4	A7 Trabzon: Çaykara, Multat plateau, alpine meadows	2811	20 vi 2018	C&Y-93
Vu5	A8 Rize: İkizdere, Ovit Lake, alpine meadows	3035	20 viii 2017	C&Y-61
<i>Vaccinium vitis-idaea</i> (Vvi)				
Vvi1	A9 Ardahan: Hanak, Alabalık plateau, alpine meadows	2362	03 vii 2018	C&Y-103
Vvi2	A9 Ardahan: Posof, Sesödile Hill, alpine meadows	2376	07 x 2017	C&Y-75
Vvi3	A8 Artvin: Şavşat Karagöl National Park, alpine meadows	2533	06 x 2017	C&Y-69
<i>Vaccinium myrtillus</i> (Vm)				
Vm1	A7 Gümüşhane: Kürtün, Süme Village, alpine meadows	1917	01 vii 2017	C&Y-5
Vm2	A8 Artvin: Şavşat, Kirazlı Village, alpine meadows	2533	02 vii 2018	C&Y-98
Vm3	A8 Rize: Ardeşen, Fındıklı Plateau, alpine meadows	2565	19 viii 2017	C&Y-52
Vm4	A9 Ardahan: Hanak, Alabalık Plateau, alpine meadows	2613	08 x 2018	C&Y-102
Vm5	A7 Trabzon: Çaykara, Multat Plateau, alpine meadows	2811	20 vi 2018	C&Y-94

^aC&Y: Coşkunçelebi and Yılmaz.

12, 14 and 17 volatile compounds were identified from the leaf of *Va*, which were collected from different altitudes in Rize-Rize-Trabzon-Trabzon-Giresun-Gümüşhane-Artvin-Artvin, respectively. Aldehydes, aliphatic hydrocarbons and aromatics were the largest group of volatiles, accounting for 44.61-77.06%, 8.59-22.01% and 3.42-23.19% of total volatiles, respectively (Table 2). In all altitudes, capronaldehyde (7.23-28.96%), 2(E)-hexenal (8.90-53.59%) and nonanal (4.24-10.03%) were generally major compounds in the leaf of *Va*.

A total of 6 to 8 volatile constituents were characterized from the leaf of *Vu*, which were harvested from different altitudes in Artvin-

Rize-Ardahan-Trabzon-Rize, respectively (Table 3). Aldehydes and aliphatic hydrocarbons were the major group of volatiles in the leaf of *Vu*, accounting for 50.92-86.95% and 8.87-30.21% of total volatiles and capronaldehyde (4.55-39.90%), 2(E)-hexenal (25.08-80.99%) and hexadecane (2.97-11.32%) were the major constituents, respectively.

The SPME analysis of *Vvi* leaf led to the identification of 6, 11 and 4 constituents and the major compounds were capronaldehyde (17.04-37.09%), α -terpinene (7.40-9.60%), limonene (16.50-47.51%) and terpinolene (7.45-8.70%), respectively (Table 4). Leaf collected from 2533 m altitude afforded limonene (47.51 %) as a major constituent

Table 2: Identified VOCs from the leaf of *V. arctostaphylos* (*Va*) growing at different altitudes in Turkey.

No	Compounds	RI*	RI ^a	<i>Va</i> 1	<i>Va</i> 2	<i>Va</i> 3	<i>Va</i> 4	<i>Va</i> 5	<i>Va</i> 6	<i>Va</i> 7	<i>Va</i> 8
				748 m	1032 m	1443 m	1516 m	1666 m	1845m	2105 m	2234 m
				(%) ^b							
1	Ethyl vinyl ketone	687	694	-	3.40	6.33	3.31	-	-	8.15	5.50
2	Pentanal	704	701	-	-	-	-	3.68	-	3.19	8.00
3	2-Ethyl furan	720	704	5.16	2.68	-	-	-	-	-	-
4	2(E)-Pentenal	756	755	-	3.46	4.78	3.49	-	-	-	-
5	Capronaldehyde	803	802	19.32	20.19	15.19	18.13	17.13	31.18	7.23	28.96
6	2(E)-Hexenal	855	854	10.4	8.90	27.50	16.26	11.01	10.70	53.59	9.30
7	Heptanal	906	903	4.17	4.38	-	-	-	-	2.23	3.78
8	Matyl caproate	925	924	-	1.55	-	-	-	-	-	-
9	2(E)-Heptenal	954	958	-	-	-	-	-	-	-	2.07
10	6-Methyl-5-hepten-2-on	981	989	9.75	3.89	-	6.66	8.76	13.50	0.96	4.06
11	2-Pentyl furan	994	994	5.30	4.30	-	5.08	7.39	4.88	3.42	5.43
12	Decane	1000	1000	-	8.30	-	-	-	-	-	-
13	Octanal	1002	1005	-	5.40	7.00	6.94	4.59	4.15	3.37	-
14	Limonene	1031	1030	-	3.02	-	0.52	-	-	-	4.46
15	2,2,6-Trimethyl cyclohexanone	1035	1041	-	-	-	-	-	-	-	3.12
16	α -Terpinene	1054	1059	-	-	-	1.96	-	-	-	-
17	(E,E)-3,5-Octadien-2-one	1068	1074	-	-	-	3.19	-	-	-	-
18	Linalool oxide	1073	1078	-	-	-	-	0.67	-	-	-
19	Linalool	1095	1102	-	-	4.36	-	2.75	3.56	-	-
20	Nonanal	1100	1106	6.79	7.58	10.03	7.14	8.26	8.53	5.81	4.24
21	2,6-Dimethylcyclohexanol	1110	1116	-	-	-	0.78	0.35	-	-	-
22	2-Allylphenol	1191	1195	6.78	3.92	11.81	5.25	15.8	9.47	-	6.97
23	Dodecane	1200	1196	-	0.92	-	-	-	-	-	-
24	Decanal	1201	1208	3.60	3.14	4.23	-	3.57	3.80	1.64	-
25	β -Cyclocitral	1224	1231	1.04	1.51	-	3.42	-	-	-	2.87
26	Vitispirane	1286	1290	5.57	-	-	-	-	-	-	-
27	Tetradecane	1400	1400	2.98	1.90	-	2.69	2.63	-	2.30	1.84
28	Geranylacetone	1453	1458	-	-	-	1.09	1.74	1.75	-	-
29	β -Ionone	1489	1496	-	-	-	1.31	-	-	-	-
30	Pentadecane	1500	1500	3.19	1.63	1.94	3.08	3.12	-	2.00	4.17
31	Dihydroactinidiolide	1537	1548	-	1.06	0.17	-	-	-	-	-
32	Hexadecane	1600	1600	5.09	3.50	3.99	5.25	5.22	4.82	3.28	2.43
33	Heptadecane	1700	1700	-	-	-	2.81	3.30	-	-	-
34	Pristane	1707	1706	8.00	4.30	2.66	1.48	-	3.65	2.76	2.73
35	Methyl myristate	1726	1728	-	1.03	-	-	-	-	-	-

36	Octadecane	1800	1800	2.85	-	-	-	-	-	-	-
	Chemical Class;										%^b and NC^c
	Monoterpene:			-	3.02:1	-	2.48:2	-	-	-	7.58:2
	Monoterpenoid:			1.04:1	1.51:1	4.36:1	3.42:1	3.42:2	3.56:1	-	-
	Aldehyde:			44.61:5	53.0:7	68.73:6	51.94:5	48.24:6	58.36:5	77.06:7	56.25:6
	Alcohol:			-	-	-	0.78:1	-	-	-	-
	Hydrocarbon:			22.01:5	20.55:6	8.59:3	16.4:6	16.01:5	10.22:3	10.34:4	11.17:4
	Aromatics:			17.14:3	14.76:3	11.81:1	10.33:2	23.19:2	14.35:2	3.42:1	12.4:2
	Ester:			-	2.58:2	-	-	-	-	-	-
	Ketone:			9.65:1	3.85:3	6.50:2	13.16:3	8.76:1	13.50:1	9.11:2	12.58:3
	Other:			5.47:1	-	-	1.31:1	-	-	-	-
	Total:			99.92:16	99.32:23	99.99:13	99.82:21	99.62:16	99.99:12	99.90:14	99.90:17

*Retention Index of references; ^aRetention Index calculated from retention times relative to that of n-alkane (C₆-C₃₂) series; ^bPercentages obtained by FID peak-area normalization; ^cNC: number of compounds; *V. arctostaphylos* species growing at different altitudes in Turkey: *Va1*: 748 m, Ardeşen-Rize; *Va2*: 1032 m Ardeşen-Rize; *Va3*: 1443 m, Araklı-Trabzon; *Va4*: 1516 m, Düzköy-Trabzon; *Va5*: 1666 m, Doğankent-Giresun; *Va6*: 1845 m Kürtün-Gümüşhane; *Va7*: 2105 m, Şavşat-Artvin; *Va8*: 2234 m, Murgul-Artvin.

Table 3: Identified VOCs and chemical class from the leaf of *V. uliginosum*(Vu) growing at different altitudes in Turkey.

No	Compounds	RI*	RI ^a	Vu1	Vu2	Vu3	Vu4	Vu5
				2530 m	2537 m	2588 m	2811 m	3035 m
				(% ^b)				
1	Pentanal	704	701	-	0.85	9.24	-	8.06
2	2-Ethyl furan	720	724	-	0.40	-	-	-
3	2(Z)-Pentenol	765	768	-	3.34	-	-	-
4	Capronaldehyde	803	802	11.67	4.55	11.18	39.90	23.16
5	4-Hydroxy-4-methyl-2-pentanone	844	845	17.47	-	20.45	-	-
6	2(E)-Hexenal	855	854	46.41	80.99	30.55	25.08	42.27
7	6-Methyl-5-hepten-2-on	981	989	-	-	-	4.81	13.27
8	Nonanal	1100	1106	4.51	-	-	-	0.48
9	Tetradecane	1400	1400	-	2.74	10.43	9.01	-
10	Pentadecane	1500	1500	-	2.56	-	-	2.37
11	Hexadecane	1600	1600	11.32	2.97	9.58	10.14	5.59
12	Heptadecane	1700	1700	8.60	-	8.55	11.06	4.78
	Chemical Class;							%^b and NC^c
	Aldehyde:			62.59:3	86.95:3	50.92:3	64.88:2	73.97:4
	Alcohol:			-	3.44:1	28.56:3	-	-
	Aliphatic hydrocarbon:			19.92:2	8.87:3	20.45:1	30.21:3	12.74:3
	Aromatics:			-	0.60:1	-	-	-
	Ketone:			17.45:1	-	-	4.81:1	13.27:1
	Total:			99.96:6	99.86:8	99.93:7	99.90:6	99.98:8

*Retention Index of references; ^aRetention Index calculated from retention times relative to that of n-alkane (C₆-C₃₂) series; ^bPercentages obtained by FID peak-area normalization; ^cNC: number of compounds; *V. uliginosum* growing at different altitudes in Turkey: *Vu1*: 2530 m, Artvin; *Vu2*: 2537 m, Rize-Anzer; *Vu3*: 2588 m, Ardahan; *Vu4*: 2811 m, Trabzon; *Vu5*: 3035 m, Rize-Ovit.

Table 4: Identified VOCs and chemical class from leaf of *V. vitis-idaea* (Vvi) growing at different altitudes in Turkey.

No	Compounds	RI*	RI ^a	Vvi1	Vvi2	Vvi3
				2362 m	2376 m	2533 m
1	Ethyl vinyl carbinol	684	689	18.13	-	-
2	Pentanal	704	701	14.60	3.90	-
3	Acetoin	711	710	-	3.05	-
4	Capronaldehyde	803	802	37.09	17.04	34.18
5	2(E)-Hexenal	855	854	9.64	-	-
6	α -Terpinene	1014	1021	-	7.40	9.60
7	Limonene	1031	1030	16.50	30.75	47.51
8	γ -Terpinene	1054	1059	-	4.60	-
9	Terpinolene	1086	1094	-	7.45	8.70
10	Linalool	1095	1102	-	6.23	-
11	Nonanal	1100	1106	4.01	1.94	-
12	α -Terpineol	1186	1192	-	14.75	-
13	Pentadecane	1500	1500	-	2.90	-
	Chemical Class;				% ^b and NC ^c	
	Monoterpene:			16.50:1	50.15:4	65.81:3
	Monoterpenoid:			-	20.98:2	-
	Aldehyde:			65.34:4	22.88:3	34.18:1
	Aliphatic hydrocarbon:			-	2.9:1	-
	Ketone:			18.13:1	-	-
	Others:			-	3.05:1	-
	Total:			99.97:6	99.96:11	99.99:4

*Retention Index of references; ^aRetention Index calculated from retention times relative to that of n-alkane (C₆-C₃₂) series; ^bPercentages obtained by FID peak-area normalization; ^cNC: number of compounds; *V. vitis-idaea* growing at different altitudes in Turkey: Vvi1: 2362 m Ardahan-Hanak; Vvi2: 2376 m Ardahan-Posof; Vvi3: 2533 m Artvin.

whereas a less percentage of limonene (16.50%) was found from other altitude (2362 m). The capronaldehyde was obtained in a high amount from the leaf collected at 2362m and 2533 m altitude. In all three sites, the monoterpene of the *Vvi* leaf collected from the high altitude was higher compared with those collected from the low altitude.

Volatile component analyzes of *Vm* leaf collected from different elevations (1917, 2533, 2565, 2613 and 2811 m) of the Black Sea region were performed by SPME GC-MS. A total of 5, 7, 12, 6 and 6 compounds were identified, respectively (Table 5). The most volatile compounds were illuminated in the sample collected from the 2565 m altitude. The main volatile constituents of *Vm* at 1917 m, 2533 m and 2565 m altitudes were capronaldehyde and 2(E)-hexenal. Aldehydes were the main class compounds, except samples harvested at 2613 m and 2811m altitude, monoterpene (39.4%) and ketone (95.27%) was the major class of compounds, respectively. The ratio of capronaldehyde and 2(E)-hexenal were decreased up to high altitudes.

DISCUSSION

In this study, the volatile component analysis of *Vaccinium* species depending on different altitudes, aldehyde compounds were found to be the main component in different proportions. In the literature, the essential oil constituents of the flowering aerial parts of *Va* were analyzed by GC-MS and twenty-six compounds, constituting 80.43% of the oil, were identified. The major compounds were α -terpineol (14.99%) and linalool (13.7%).⁸ The volatile oil from *V. bracteatum* leaf was analyzed

by GC-MS and 49 constituents were reported, which represented about 91.94% of the total content. The main chemical compounds were terpenes (54.75%), open chain alkanes (6.93%), cycloparaffins (20.1%), olefins (0.60%), aromatics (1.60%), phenol, ether and alcohol (3.39%), aldehyde and ketone (2.14%), ester (1.15%), heterocyclic compounds (0.47%) and amines (0.78%). Nerolidol (20.01%), (Z,Z,Z)-1,5,9,9-tetramethyl-1,4,7-cycloundecatriene (17.99%) and caryophyllene (9.59%) were the main components in the identified terpenes.⁹ The GC/MS analyses of the essential oils from dry leaf *Vv i* were predominant to be terpenoids, fatty acids, fatty acid- and carotenoid-derived compounds and α -terpineol (4.7%) and linalool (17.0%).¹⁰ The chemical constituents of blueberry seed oil was analyzed by GC-MS and a total of 34 constituents were reported. The contents in the volatiles were menthyl isovalerate (30.06%), 9,12-octadecadien-1-ol (11.82%), Z-7-tetradecenol (9.77%), γ -sitosterol (6.04%), β -sitosterol (4.74%), phenylethyl alcohol (4.72%), Z-(13,14-epoxy)tetradec-11-en-1-ol acetate (4.30%), retinal (4.07%), 1-(+)-ascorbic acid, 6-dihexadecanoate (3.93%), 9,12-octadecadienoic acid, ethyl ester (3.10%) and 1,6,10,14,18,22-tetracosahexaen-3-ol (3.00%).³² The chemical compounds of the essential oils obtained from by hydrodistillation from the flowering aerial parts of *Va* was assessed by GC/MS analysis. α -Pinene (15.5%), linalool (11.7%), sandaracopimaradiene (5.9%) and safranal (8.8%) were mentioned to be major constituents.¹¹ The volatile components of *Vvi* juice were analyzed by HS SPME GC-MS and 41 volatile components were detected. The major volatile compound in *Vvi* juice was reported as diisobutyl phthalate, accounting for 17.96% of total aromatic

Table 5: Identified VOCs and chemical class from leaf of *V. myrtillus* (Vm) growing at different altitudes in Turkey.

No	Compounds	RI*	RI ^a	Vm1	Vm2	Vm3	Vm4	Vm5
				1912 m	2533 m	2565 m	2613 m	2811 m
1	Ethyl vinyl carbinol	684	689	-	-	-	-	4.22
2	3-Penten-2-one	733	739	9.44	-	14.52	93.60	-
3	Capronaldehyde	803	802	20.26	11.50	37.26	1.01	14.66
4	4-Hydroxy-4-methyl-2-pentanone	844	845	19.94	-	-	1.67	-
5	2(E)-Hexenal	855	854	37.59	15.00	18.18	1.31	34.13
6	α -Pinene	932	938	-	12.55	-	-	-
7	1-Octen-3-ol	976	979	-	-	-	-	2.70
8	6-Methyl-5-hepten-2-on	981	989	3.86	-	13.82	-	9.36
9	2-Pentyl furan	994	994	-	-	-	-	1.65
10	<i>p</i> -Cymene	1020	1029	-	7.12	-	-	-
11	Limonene	1031	1030	-	19.99	-	-	-
12	2,2,6-trimethyl Cyclohexanone	1035	1041	-	-	-	-	4.94
13	Nonanal	1100	1106	8.88	-	2.87	-	2.58
14	2-Bornanone	1161	1154	-	32.86	-	-	-
15	β -Cyclocitral	1224	1231	-	-	-	0.71	2.76
16	Vitispirane	1286	1290	-	-	-	1.30	-
17	Tridecane	1300	1299	-	-	-	-	6.60
18	Tetradecane	1400	1400	-	-	-	-	8.03
19	Hexadecane	1600	1600	-	-	8.70	-	4.57
20	Heptadecane	1700	1700	-	-	4.61	-	3.75
	Chemical Class;					% ^b and NC ^c		
	Monoterpene:			-	39.4:3	-	-	-
	Monoterpenoid:			-	32.9:1	-	0.71:1	2.76:1
	Aldehyde:			66.73:3	27.6:2	58.31:3	2.32:2	51.37:3
	Alcohol			-	-	-	-	2.70:1
	Aliphatic hydrocarbon:			-	-	13.31:2	-	22.95:4
	Ketone:			33.24:3	-	28.34:2	95.27:2	18.52:3
	Others:			-	-	3.05:1	1.3:1	-
	Total:			99.97:6	99.99:6	99.96:7	99.96:6	99.95:12

*Retention Index of references; ^aRetention Index calculated from retention times relative to that of n-alkane (C₆-C₃₂) series; ^bPercentages obtained by FID peak-area normalization; ^cNC: number of compounds; *V. myrtillus* species growing at different altitudes in Turkey: Vm1: 1917m, Gümüşhane-Kürtün; Vm2: 2533m, Artvin-Şavşat; Vm3: 2565m, Rize-Ardeşen; Vm4: 2613m, Ardahan-Hanak; Vm5: 2811m, Trabzon-Çaykara.

compounds.¹² The volatile components from the fresh leaf buds of *V. dunalianum* var. *dunalianum* were mentioned. 27 Compounds has been identified and decahydro dimethylnaphthalene (47.04%), linalool (3.19%), 3,7-dimethyl-1,5,7-octatrien-3-ol (1.40%) and benzene acetaldehyde (1.03%) were the main compounds.¹³ The floral volatile compounds of *V. angustifolium*, *V. varingiaefolium*, *V. arboreum* and *V. poasanum* has been provided. A total of 40 from *V. angustifolium*, 34 for *V. varingiaefolium*, 37 for *V. arboreum* and 17 for *V. poasanum* volatile compounds were detected in the headspace surrounding flowers. The most abundant volatile compounds for the each species was reported to be α -pinene for *V. angustifolium* (23%) and *V. arboreum* (63%), methyl benzoate for *V. varingiaefolium* (18%) and ethyl benzene for *V. poasanum* (21%).¹⁴ Five volatile compounds consisting of butyl butanoate, *cis*-3-hexen-1-ol, α -terpineol, geraniol and 2(E)-hexenal has been found from *V. corymbosum* using GC-MS.¹⁵ Vm juice has been analyzed by HS SPME GC-MS and altogether, 56 volatile compounds were identified.¹⁶

The complexity of Vm aroma has been explored by SPME GC-MS. This comprehensive characterization of aroma allowed the identification of volatile organic compounds, for the most aldehydes, alcohols, terpenoids, and esters.¹⁷ A study has been conducted to determine the antifungal and the phytotoxic activity of Vm plant essential oil.¹⁸ In another work, volatile compounds and the antifungal activity of Bilberry essential oil of Vm were mentioned and 22 components were identified as mainly 1,8-cineole (41.07%), β -linalool (12.72%), α -pinene (12.17%) and myrtenol (6.48%).¹⁹ Polar metabolites of leaf of Vm and Vvi has been studied by HPLC-DAD, HPLC-MS and GC-FID. Chlorogenic acid was the major phenolic compounds in bilberry leaf and arbutin in lingonberry leaf. Flavonol glycosides were another major group of phenolic in bilberry [5-28 mg/g DM (dry mass)] and lingonberry (15-20 mg/g DM) leaf.²⁰ The extracts of entire fruits and leaf of Vm collected in Finland and Poland has been quantitated by GC-FID/MS and the

main bilberry constituents were found to be α - and β -amyrin, α - and β -amyrenone, campesterol, cholesterol and citrostadienol.²¹

CONCLUSION

Vaccinium species is one of the industrial medicinal plant that is traditionally used for the treatment of diabetes throughout the world. VOCs composition of the *Va*, *Vu*, *Vvi* and *Vm* leaf has been analyzed. Comparisons based on the mean relative amount of the volatile components in different sites showed significant differences among populations at different altitudes. In all three and five sites, the monoterpene and aldehydes ratio of the *Vvi*, *Vm* and *Vu* collected from the low altitude was higher compared with those collected from the high altitude, respectively. The aldehydes ratio of the *Va* harvested from the all eight altitudes was more or less the same. Principal components analyzed based on the mean relative amounts of the volatile components have led to the identification of three chemotypes (capronaldehyde, 2(*E*)-hexenal and nonanal). Since, the amount of plant constituents affected by altitude and time collected would vary accordingly. These results suggest that altitude is a factor influencing the volatile components of these plants.

ACKNOWLEDGEMENT

This research was supported by KTÜ-BAP (TYL-2017-6714) and the authors are thankful to the Faculty of Pharmacy research facilities, Karadeniz Technical University, Turkey.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ABBREVIATIONS

Va: *V. arctostaphylos*; **Vu:** *V. uliginosum*; **Vvi:** *V. vitis-idaea*; **Vm:** *V. myrtillus*; **SPME:** Solid phase micro extraction; **GC-MS:** Gas Chromatography-Mass Spectrometry; **HS:** Headspace; **VOC:** Volatile organic compound; **HPLC:** High-Performance Liquid Chromatography; **DVB:** Divinylbenzene; **PDMS:** Polydimethylsiloxane; **FID:** Flame ionization detector; **mL:** Millilitre; **KTUB:** Karadeniz Technical University, Herbarium of Biology (Turkey); **C&Y:** Coşkunçelebi and Yılmaz; **RI:** Retention Index; **NC:** Number of compounds.

REFERENCES

- Cronquist A. An integrated system of classification of flowering plants. New York: Columbia University Press. 1981;1262.
- Stevens PFA. Classification of the Ericaceae: Subfamilies and tribes. Bot J Linn Soc. 1971;64(1):1-53.
- Vander KSP. The genus *Vaccinium* in North America. Research Branch Publish Agriculture: Canada, Ottawa. 1988.
- Martineau LC, Couture A, Spoorab D, Benhaddou-Andaloussi A, Harris C, Meddahab B, et al. Anti-diabetic properties of the Canadian lowbush blueberry *Vaccinium angustifolium* Ait. Phytomedicine. 2006;13(10):612-23.
- Bljajić K, Petlevski R, Vujić L, Čačić A, Šoštarčić N, Jablan J, et al. Chemical composition, antioxidant and α -glucosidase-inhibiting activities of the aqueous and hydroethanolic extracts of *Vaccinium myrtillus* leaves. Molecules. 2017;22(5):703.
- Ayaz FA, Hayirlioğlu-Ayaz S, Gruz J, Novak O, Strnad M. Separation, characterization and quantitation of phenolic acids in a little-known blueberry (*Vaccinium arctostaphylos* L.) fruit by HPLC-MS. J Agric Food Chem. 2005;53(21):8116-22.
- Rimando AM, Kalt W, Magee JB, Dewey J, Ballington JR. Resveratrol, pterostilbene and piceatannol in *Vaccinium* berries. J Agric Food Chem. 2004;52(15):4713-9.
- Nickavar B, Salehi-Sormagi M, Amin G, Daneshlab M. Steam volatiles of *Vaccinium arctostaphylos*. Pharm Biol. 2002;40(6):448-9.
- Yang X, Xiao S, Xu Y, Han Z, Wu Y. Analysis of volatile oil from *Vaccinium bracteatum* Thumb. leaves by gas chromatography-mass spectrometry. Biomass Chem Eng. 2008;42(2):23-6.
- Radulovic N, Blagojevic P, Palic R. Comparative study of the leaf volatiles of *Arctostaphylos uva-ursi* (L.) Spreng. and *Vaccinium vitis-idaea* L. (Ericaceae). Molecules. 2010;15(9):6168-85.
- Teimouri M. The chemical composition and antimicrobial activity of essential oils of *Vaccinium arctostaphylos* L. Int J Adv Biol Biomed Res. 2014;2(11):2837-41.
- Yang H, Liu Y, Guo D. Volatile components analysis of *Vaccinium vitis-idaea* juice and its fermented wine by GC-MS. Zhongguo Niangzao. 2014;33(12):133-8.
- Yin J, Jiang L, Jiang Y, Wang J, Zhao P. GC-MS analysis of volatile components from fresh leaf-buds of *Vaccinium dunalianum* var. *dunalianum*. J Southwest Forestry Univer. 2013;33(2):100-3.
- Forney CF, Javorek SK, Jordan MA, Vander KSP. Floral volatile composition of four species of *Vaccinium*. Botany. 2012;90(5):365-71.
- Liburd OE. Identification of host volatile compounds for monitoring blueberry maggot fly. Int J Fruit Sci. 2004;3(3/4):307-12.
- Liu S, Laaksonen O, Yang B. Volatile composition of bilberry wines fermented with non-Saccharomyces and Saccharomyces yeasts in pure, sequential and simultaneous inoculations. Food Microbiol. 2019;80:25-39.
- Farneti B, Grisenti M, Ajelli M, Giongo L, Khomenko I, Betta E, et al. Exploring blueberry aroma complexity by chromatographic and direct-injection spectrometric techniques. Front Plant Sci. 2017;8:617.
- Yılar M. Phytotoxic and antifungal activities of thyme, bilberry, sage essential oils. Fresenius Environ Bull. 2018;27(8):5559-69.
- Bayar Y, Onaran A, Yılar M, Gul F. Determination of the essential oil composition and the antifungal activities of bilberry (*Vaccinium myrtillus* L.) and bay laurel (*Laurus nobilis* L.). J Essent Oil Bear Pl. 2018;21(2):548-55.
- Liu P, Lindstedt A, Markkinen N, Sinkkonen J, Suomela J, Yang B. Characterization of metabolite profiles of leaves of bilberry (*Vaccinium myrtillus* L.) and lingonberry (*Vaccinium vitis-idaea* L.). J Agric Food Chem. 2014;62(49):12015-26.
- Szakiel A, Paczkowski C, Huttunen S. Triterpenoid content of berries and leaves of bilberry *Vaccinium myrtillus* from Finland and Poland. J Agric Food Chem. 2012;60(48):11839-49.
- Renda G, Tosun G, Yaylı N. SPME GC/MS analysis of three *Ornithogalum* L. species from Turkey. Rec Nat Prod. 2016;10(4):497-502.
- Korkmaz FM, Özel MB, Tüzüner T, Korkmaz B, Yaylı N. Antimicrobial activity and volatile constituent analysis of three commercial herbal toothpastes containing *Aloe vera* L. and *Fragaria vesca* L. extracts. Niger J Clin Pract. 2019;22(5):718726.
- Renda G, Çelik G, Korkmaz B, Alpay Ş, Yaylı N. Antimicrobial activity and analyses of six *Geranium* L. species with headspace SPME and hydrodistillation. J Essent Oil Bear Pl. 2016;19(8):2003-16.
- Shelke V, Bhot M. GC-MS Analysis of bio-active compounds in ethanolic extract of leaf and stem of *Asclepias curassavica* L. Int J Pharm Investig. 2019;9(2):67-70.
- Adams RP. Identification of essential oil components by gas chromatography/mass spectrometry. Allured Publishing Corporation Carol Stream I. 2007.
- Adams RP, Morris JA, Pandey RN, Schwarzbach AE. Cryptic speciation between *Juniperus deltoides* and *Juniperus oxycedrus* (Cupressaceae) in the mediterranean. Biochem Syst Ecol. 2005;33(8):771-87.
- Yue T, Chi M, Song C, Liu M, Jiang-Fei Y, Zhang Z, et al. Aroma characterization of cabernet sauvignon wine from the plateau of yunnan (China) with different altitudes using SPME-GC/MS. Int J Food Prop. 2015;18(7):1584-96.
- Bylaite E, Meyer AS. Characterisation of volatile aroma compounds of orange juices by three dynamic and static headspace gas chromatography techniques. Eur Food Res Technol. 2006;222(1-2):176-84.
- Nickavar B, Salehi-Sormagi MH, Amin G, Daneshlab M. Steam volatiles of *Vaccinium arctostaphylos*. Pharm Biol. 2002;40(6):44849.
- Quijano CE, Salamanca G, Pino JA. Aroma volatile constituents of Colombian varieties of mango (*Mangifera indica* L.). Flavour Frag J. 2007;22(5):401-16.
- Qu X, Bai Y, Li Y, Yang Y, Jia Q, Nie F. Analysis on volatiles from blueberry seed oil by GC-MS. Guizhou Nongye Kexue. 2013;41(10):6466.

Article History: Submission Date : 25-08-2020; Revised Date : 15-09-2020; Acceptance Date : 20-10-2020.

Cite this article: Erik İ, Kılıç G, Yaylı SS, Yılmaz G, Coşkunçelebi K, Yaylı N. Altitude Variation in Volatile Composition of Blueberry Leaf Analyzed by SPME GC-MS. Int. J. Pharm. Investigation, 2020;10(4):519-25.