

Antibiotics Removal from Aqueous Solution and Pharmaceutical Wastewater by Adsorption Process: A Review

Davoud Balarak¹, Aram Dokht Khatibi¹, Kethineni Chandrika^{2*}

¹Department of Environmental Health, Health Promotion Research Center, Zahedan University of Medical Sciences, Zahedan, IRAN.

²Department of Biotechnology, Koneru Lakshmaiah Education Foundation, Vaddeswaram, Guntur, Andhra Pradesh, INDIA.

ABSTRACT

With the widespread use of antibiotics, pharmaceutical effluents containing antibiotics have recently attracted wide attentions since it has potential adverse effects. Although the injuriousness of antibiotics is not so intuitive like other environmental pollution, residues of the antibiotic drug have become a seriously ignored problem. The abuse of antibiotics can damage the immune function of animals, when again infected, and then they need more antibiotics to treat. In this review article the authors presented up to-date development on the application of adsorption in the removal of antibiotics from aqueous solution. This review article provides extensive literature information about antibiotics, its classification and toxicity, various treatment methods, and antibiotics adsorption characteristics by various adsorbents. The effectiveness of various adsorbents under different physico-chemical process parameters and their comparative adsorption capacity towards antibiotics adsorption has also been presented. This review paper

also includes the affective adsorption factors of antibiotics such as solution contact time, initial antibiotics concentration, adsorbent dosage, and temperature. The applicability of various adsorption thermodynamic models and isotherm models for antibiotics removal by wide range of adsorbents is also reported here.

Key words: Adsorbent, Antibiotics, Adsorption, Wastewater, Thermodynamic.

Correspondence

Prof. Kethineni Chandrika,

Associate Professor, Department of Biotechnology, Koneru Lakshmaiah Education Foundation, Vaddeswaram, Guntur, Andhra Pradesh, INDIA.

Email: kkchandrka2019@gmail.com

DOI: 10.5530/ijpi.2020.2.19

INTRODUCTION

Antibiotics are widely used in the treatment of diseases in animals and humans, as well as applied in food rations in order to increase animal growth rates.¹⁻³ Antibiotics have high antimicrobial activity associated it's the aromatic structure containing the naphthol chemical group that is antibacterial.⁴ However, approximately 50-90% of the doses of antibiotics administered in health treatment procedures is not absorbed by organisms and is eliminated by humans and animals in sewage systems.⁵ These molecules have a complex chemical structure and the natural environmental and sewage conditions are not sufficient to decompose their chemical structure.⁶ Thus, the amount of antibiotics accumulated in sewage can be a serious environmental problem.⁷ Due to the chemical structure of antibiotics, they are act as a resistant to many chemicals, oxidizing agents, and heat, and are biologically non-degradable.⁸ So it is difficult to removal the effluents, once released into the aquatic environment.⁹ Many of the methods are available for the removal of pollutants from water, the most important of which are reverse osmosis, ion exchange, precipitation and adsorption.¹⁰

Adsorption is used as top quality treatment procedures for the removal of dissolved organic pollutants like antibiotics from industrial waste water.¹¹ Adsorption is defined as concentration of materials on the surface of solid bodies.¹² Adsorption is a surface phenomenon which deals primarily with the utilization of surface forces.¹³ When a solution having absorbable solute, also called as adsorbate, comes into contact with a solid, called as adsorbent, with highly porous surface structure liquid-solid intermolecular forces of attraction causes the solute to be concentrated at the solid surface.¹⁴ Adsorption is one of the unit operations in the chemical engineering processes used for the separation of industrial wastewater pollutants.¹⁵

The present state of the art on the application of adsorption in the removal of antibiotics from aqueous solution is presented in this paper. The main goal of this review article is to provide up-to-date development on the application of commercial activated carbon and various sustainable low cost alternative adsorbents such as agricultural solid waste (azolla, Lemna minor, canola and ...) , industrial solid waste, agricultural by-products, and biomass based cost effective activated carbon, and various natural materials in the removal of antibiotics from aqueous phase. This review article also critically analyses the effectiveness of various adsorbents under different physicochemical process parameters and their comparative adsorption capacity is also presented. A compilation of relevant published data with respect to adsorption kinetics, isotherm models, thermodynamics and adsorption capacity under various process conditions along with important findings is presented here.

Although there are a couple of review articles such as low cost adsorbents for the removal of organic pollutants from wastewater,¹⁶⁻¹⁸ but all are dealing with specific system only and also not relatively presented with up-to date information. Therefore, this present review article was undertaken in order to provide more comprehensive up-to-date and critical review information on the adsorption of various antibiotics from aqueous solution by wide range of adsorbents. The new aspect of this review article is to cover up-to-date research result presentation on various antibiotics adsorption and its adsorptive effectiveness in the removal of various antibiotics and also to critically analyzed and identify various operation conditions and their maximum adsorption capacity. Authors also tried to analyze the scattered available information on antibiotics adsorption by wide range of adsorbents since the last two decades.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

METHODOLOGY

This systematic review aimed to study the antibiotics adsorption from aqueous solutions using different adsorbent. Studies published in electronic form in international database of Google Scholar, Scopus, PubMed, Web of Science and Science Direct from 2000 until end of the 2019 year, was investigated. Search using antibiotic, adsorption, efficacy and removal keywords with AND/OR operators in title and abstract was conducted.

RESULTS

Table 1 shows the reported studies on the effect of adsorbent dosage on the percentage of antibiotics removal. Table 2 showed the compilation of results of various studies on the effect of temperature on the antibiotics

adsorption by various adsorbents. Table 3 shows the various isotherm results of antibiotics adsorption by various adsorbents. Table 4 also shows the maximum adsorption capacity (q_m) of different adsorbents in the field of antibiotic removal obtained from the Langmuir equation.

DISCUSSION

The term adsorption refers to the accumulation of a substance at the interface between two phases (liquid–solid interface or gas–solid interface).¹⁰³ The substance that accumulates at the interface is called adsorbate and the solid on which adsorption occurs is adsorbent.¹⁰⁴ Adsorption can be classified into two types: chemical sorption and physical sorption. Chemical adsorption or chemisorption is illustrated by the formation of strong chemical associations between molecules or ions of adsorbate to adsorbent surface, which is generally due to

Table 1: The results of various reported studies on the effect of adsorbent dose on antibiotics adsorption by various adsorbents.

Adsorbents	Antibiotics name	Adsorbent dosage	%Removal	References
Carbon materials	Amoxicillin	0.05-0.5 g/L	39.5-75.6	19
AC-vine wood	Cephalexin	0.1-1 g/L	44.5-87.2	20
Bentonite	Amoxicillin	0.25-2.5 g/L	41.8-94.6	21
AC-siris seed	Metronidazole	0.15-0.8 g/L	35.2- 64.2	22
AC	tetracycline	0.05-0.6 g/L	44.6-78.9	23
Montmorillonite	Penicillin G	0.4-2 g/L	64.8-98.2	24
AC-lignocellulosic	Cephalexin	0.15-0.7 g/L	21.9-92.1	25
NiO nanoparticle	Amoxicillin	0.1-0.5 g/L	54.3-87.9	26
AC- pyrolysis char	Tetracycline	0.02-1 g/L	71.5-91.8	27
AC commercial	Ibuprofen	0.25-1.5 g/L	61.4-97.4	28
Maize stalks	Tetracycline	0.5-4 g/L	54.6-94.2	29
Azolla filiculoides	Tetracycline	0.5-5 g/L	69.8-87.9	30
AC-macadamia shells	Tetracycline	0.3-1.5 g/L	71.4-87.2	31
Granular sludge	Oxytetracycline	0.4-2 g/L	39.8-79.8	32
Chestnut shell	penicillin G	0.5-3 g/L	28.7-81.4	33

Table 2: The effect of temperature on the adsorption of Antibiotics using various adsorbents.

Adsorbents	Antibiotics name	Temperature (K)	Results	References
Grapheme oxide magnetic	Tetracycline	273-323	Endothermic	34
Activated alumina	Metronidazole	273-303	Exothermic	35
Rice husk	Metronidazole	273-303	Endothermic	36
MWCNT	Metronidazole	288-318	Endothermic	37
Azolla Filiculoides	Cephalexin	293-333	Endothermic	38
mesoporous silica	Tetracycline	273-313	Exothermic	39
Cu-13X	Tetracycline	288-318	Endothermic	40
MWCNT	Tetracycline	283-313	Exothermic	41
Zero-valent iron	Tetracycline	273-313	Endothermic	42
AC-apricot nut shells	Tetracycline	273-323	Endothermic	43
AC commercial	Quinolone	283-313	Endothermic	44
NiO nanoparticle	Amoxicillin	288-318	Exothermic	45
CuO nanoparticles	Ciprofloxacin	288-323	Endothermic	46
MWCNT	Amoxicillin	273-323	Endothermic	47
Graphene oxide	Tetracycline	283-313	Exothermic	48

Table 3: Various isotherm studies of antibiotics adsorption by various adsorbents.

Adsorbents	Antibiotics name	studied isotherms	best isotherm	References
Chitosan	Amoxicillin	Langmuir-Freundlich-Temkin-D-R	Langmuir	49
Composite iron nano	Ibuprofen	Langmuir-Freundlich-Temkin-R-P	D-R	50
Carbon xerogel	Ciprofloxacin	Langmuir-Freundlich-BET-D-R	Langmuir	51
Polyoxometalates-NP	Tetracycline	Langmuir-Freundlich-Temkin-BET	BET	52
Fe ₃ O ₄ -graphene	Tetracycline	Langmuir-Freundlich-BET-D-R	D-R	53
Hydroxyapatite	Tetracycline	Langmuir-Freundlich-Temkin- D-R	D-R	54
Magnetic-MWCNT	Amoxicillin	Langmuir-Freundlich-Temkin-BET	Langmuir	55
Rice husk ash	Tetracycline	Langmuir-Freundlich-Temkin-BET	Temkin	56
MWCNT	Cephalosporins	Langmuir-Freundlich-Temkin-D-R	Langmuir	57
MWCNT	Tetracycline	Langmuir-Freundlich-Temkin-BET	Temkin	58
SWCNT	Ibuprofen	Langmuir-Freundlich-Temkin-R-P	Freundlich	59
Granular AC	Trimethoprim	Langmuir-Freundlich-Temkin- D-R	Langmuir	60
Coal fly ash	Norfloxacin	Langmuir-Freundlich-Temkin	Langmuir	61
Coal fly ash	Ciprofloxacin	Langmuir-Freundlich-Temkin- D-R	Langmuir	62
Activated sludge	Tetracycline	Langmuir-Freundlich-Temkin	Temkin	63
Canola	Amoxicillin	Langmuir-Freundlich-Temkin-R-P	Freundlich	64
Montmorillonite	Tetracycline	Langmuir-Freundlich-Temkin-BET	Langmuir	65

Table 4: Adsorption capacity of different adsorbents in the field of antibiotic removal.

Adsorbents	Antibiotics	Q _m (mg/g)	Ref	Adsorbents	Antibiotics	Q _m (mg/g)	Ref
NH ₄ Cl-AC	Amoxicillin	28.24	66	Iron particles	Ampicillin	41.17	67
Azolla filiculoides	Ampicillin	19.45	68	Aluminum oxide	Tetracycline	71.49	69
Fe-Mn oxide	Tetracycline	82.57	70	Goethite	Tetracycline	35.79	71
Activated sludge	Tetracycline	41.43	72	Chitosan	Tetracycline	28.64	73
MnFe ₂ O ₄ /AC	Tetracycline	73.96	74	Titania-silica	Tetracycline	41.72	75
Azolla -AC	Amoxicillin	98.47	76	Magnetic-CNT	Amoxicillin	148.2	77
Lemna minor	Penicillin G	29.47	78	Montmorillonite	Tetracycline	36.94	79
Kaolinite	Tetracycline	41.56	80	Illite	Tetracycline	38.74	81
Cay minerals	Tetracycline	35.68	82	Lemna minor algal	Tetracycline	36.47	83
Wheatstalks	Tetracycline	21.79	84	Corylus -AC	Ciprofloxacin	125.9	85
Azolla-AC	Sulfamethoxazole	119.7	86	Alumina-CNT	Tetracycline	264.8	87
Graphite	Tetracycline	84.35	88	Grapheme oxide	Tetracycline	114.6	89
Bntonite	Ampicillin	61.74	92	Palygorskite	Tetracycline	146.5	91
Silica	Tetracycline	48.14	92	Polyaniline-NP	Metronidazole	36.94	93
Zirconium Oxide-NP	Ciprofloxacin	219.8	94	Zeolite A	Clinoptilolite	181.2	95
siris seed pods- AC	metronidazole	81.46	96	NH ₄ Cl- AC	Tetracycline	92.17	97
Ativated carbons	Tetracycline	95.27	98	SiO ₂ -NP	Ciprofloxacin	84.91	99
Red mud	Ciprofloxacin	52.14	100	Plm bark	Amoxicillin	29.65	101
MAC	Ciprofloxacin	156.4	102	Cu-13X	Tetracycline	64.81	40
Fe ₃ O ₄ -graphene	Tetracycline	54.62	53	Canola	Amoxicillin	24.19	64

the exchange of electrons and thus chemical sorption generally is irreversible.¹⁰⁵ Physical adsorption or physisorption is characterized by weak Van der Waals intraparticle bonds between adsorbate and adsorbent and thus reversible in most cases²². Adsorption on most of the adsorbent including agricultural by-products is controlled by physical forces with some exception of chemisorption.¹⁰⁶ The main physical forces controlling adsorption are Van der Waals forces, hydrogen bonds,

polarity, dipole-dipole $\Pi-\Pi$ interaction, etc.¹⁰⁷ This process provides an attractive alternative for the treatment of polluted waters, especially if the sorbent is inexpensive and does not require an additional pretreatment step before its application.²⁴ As for environmental remediation purpose, adsorption techniques are widely used to remove certain classes of chemical contaminants from waters, especially those that are practically unaffected by conventional biological wastewater treatments.²⁵ Adsorption

has been found to be superior to other techniques in terms of flexibility and simplicity of design, initial cost, insensitivity to toxic pollutants and ease of operation.²⁶ Adsorption also does not produce harmful substances.²⁸ Factors that influence the adsorption efficiency include adsorbate-adsorbent interaction, adsorbent surface area, adsorbent to adsorbate ratio, adsorbent particle size, temperature, pH and contact time and etc. Adsorbent dosage is an important process parameter to determine the capacity of an adsorbent for a given amount of the adsorbent at the operating conditions.¹⁰⁸ Generally the percentage of antibiotics removal increases with increasing adsorbent dosage, where the quantity of sorption sites at the surface of adsorbent will increase by increasing the amount of the adsorbent.¹⁰⁹ The effect of adsorbent dosage gives an idea for the ability of a antibiotics adsorption to be adsorbed with the smallest amount of adsorbent, so as to recognize the capability of a antibiotics from an economical point of view.

Effect of temperature is another significant physico-chemical process parameter because temperature will change the adsorption capacity of the adsorbent.¹⁰⁸ If the amount of adsorption increases with increasing temperature then the adsorption is an endothermic process.¹⁰⁹ This may be due to increasing mobility of the antibiotics molecules and an increase in the number of active sites for the adsorption with increasing temperature.¹¹⁰ Whereas the decrease of adsorption capacity with increasing temperature indicates that the adsorption is an exothermic process.¹¹¹ This may be due to increasing temperature decreasing the adsorptive forces between the antibiotics species and the active sites on the adsorbent surface as a result of decreasing the amount of adsorption.¹¹² Table 2 showed the compilation of results of various studies on the effect of temperature on the antibiotics adsorption by various adsorbents.

The adsorption isotherm is significant for the explanation of how the adsorbent will interact with the adsorbate and give an idea of adsorption capacity. They play an important role in understanding the mechanism of adsorption. The surface phase may be considered as a monolayer or multilayer. Several isotherm models are presented in the literature.¹¹³ Langmuir and Freundlich models are the most widely used to describe the adsorption isotherm. The Langmuir adsorption isotherm model assumed that adsorption takes place at specific homogeneous sites within the adsorbent, and it has been used successfully for many adsorption processes of monolayer adsorption.¹¹⁴ The Freundlich adsorption isotherm model considers a heterogeneous adsorption surface that has unequal available sites with different energies of adsorption.¹¹⁵

CONCLUSION

In review article presented a wide range of adsorbents such as nanoparticles, carbon nano tubes, various agricultural adsorbent, activated carbon, biomass-based activated carbon, biosorbents, various other inorganic oxides, and clay minerals in the removal of antibiotics from aqueous solution. From the large number of published literature reviewed here, it is observed that the mechanism and kinetics of adsorption of antibiotics on various adsorbents depend on the chemical nature of the materials and various physico-chemical experimental conditions such as initial antibiotics concentration, adsorbent dosage, and temperature of the system. Therefore, these factors are to be taken into account while evaluating the adsorption capacity of different adsorbents. This review article also reveals that the Langmuir and Freundlich adsorption isotherm models are usually used to evaluate the adsorption capacity of various adsorbents.

REFERENCES

- Klavarioti M, Mantzavinos D, Kassinos D. Removal of residual pharmaceuticals from aqueous systems by advanced oxidation processes. *Environ Int.* 2009;35(2):402-17.
- Ahmadi M, Motlagh HR, Jaafarzadeh N, Mostoufi A, Saeedi R, Barzegar G, *et al*. Enhanced photocatalytic degradation of tetracycline and real pharmaceutical wastewater using MWCNT/TiO₂ nano-composite. *J Environ Manage.* 2017;186(1):55-63.
- Huang CH. Assessment of potential antibiotic contaminants in water and preliminary occurrence analysis. *J Contemp Water Res Educ.* 2011;120(1):4-11.
- Álvarez-Torrellas S, Rodríguez A, Ovejero G, García J. Comparative adsorption performance of ibuprofen and tetracycline from aqueous solution by carbonaceous materials. *Chem Eng J.* 2016;283:936-47.
- Chayid MA, Ahmed MJ. Amoxicillin adsorption on microwave prepared activated carbon from arundodonaxlinn: Isotherms, kinetics, and thermodynamics studies. *J Environ Chem Eng.* 2015;3(3):1592-601.
- Carabineiro SAC, Thavorn-Amornsri T, Pereira MFR, Figueiredo JL. Adsorption of ciprofloxacin on surface-modified carbon materials. *Water Res.* 2011;45(15):4583-91.
- Chingombe P, Saha B, Wakeman RJ. Sorption of atrazine on conventional and surface modified activated carbons. *J Colloid Interf Sci.* 2014;302(2):408-16.
- Choi KJ, Kim SG, Kim SH. Removal of antibiotics by coagulation and granular activated carbon filtration. *J Hazard Mater.* 2008;151(1):38-43.
- Conkle JL, Lattao C, White JR, Cook RL. Competitive sorption and desorption behavior for three fluoroquinolone antibiotics in a waste water treatment wetland soil. *Chemosphere* 2010;80(11):1353-9.
- Shafey EI, Al-Lawati H, Al-Sumri AS. Ciprofloxacin adsorption from aqueous solution onto chemically prepared carbon from date palm leaflets. *J Environ Sci.* 2012;24(9):1579-86.
- Laak TL, Gebbink VA, Tolls J. The effect of pH and ionic strength on the sorption of sulfachloropyridazine, tylosin, and oxytetracycline to soil. *Environ Toxicol Chem.* 2006;25(4):904-11.
- Zhang L. Studies on the removal of tetracycline by multi-walled carbon nanotubes. *Biochem Eng J.* 2011;178:26-33.
- Zhang X. Adsorption of sulfamethoxazole on different types of carbon nanotubes in comparison to other natural adsorbents. *J Environ Sci health part A.* 2010;45(12):1625-34.
- Zhang D, Yin J, Zhao J, Zhu H, Wang C. Adsorption and removal of tetracycline from water by petroleum coke-derived highly porous activated carbon. *J Environ Chem Eng.* 2015;3(3):1504-12.
- Zhu X, Liu Y, Qian F, Zhou C, Zhang S, Chen J. Preparation of magnetic porous carbon from waste hydrochar by simultaneous activation and magnetization for tetracycline removal. *Bioresour Technol.* 2014;154:209-14.
- Ocampo-Pérez R, Rivera-Utrilla J, Gómez-Pacheco C, Sánchez-Polo M, López-Peñalver JJ. Kinetic study of tetracycline adsorption on sludge-derived adsorbents in aqueous phase. *Chem Eng J.* 2012;213:88-96.
- Sun Y, Yue Q, Gao B, Huang L, Xu X, Li Q. Comparative study on characterization and adsorption properties of activated carbons with H₃PO₄ and H₄P₂O₇ activation employing *Cyperus alternifolius* as precursor. *Chem Eng J.* 2012;181:790-7.
- Suga N, Ogo M, Suzuki S. Risk assessment of oxytetracycline in water phase to major sediment bacterial community: a water-sediment microcosm study. *Environ Toxicol Pharmacol.* 2013;36(1):142-8.
- Yu F, Li Y, Han S, Ma J. Adsorptive removal of antibiotics from aqueous solution using carbon materials. *Chemosphere.* 2016;153:365-85.
- Pouretedal HR, Sadegh N. Effective removal of amoxicillin, cephalixin, tetracycline and penicillin G from aqueous solutions using activated carbon nanoparticles prepared from vine wood. *J Water Process Eng.* 2014;1:64-73.
- Putra EK, Pranowo R, Sunarso J, Indraswati N, Ismadji S. Performance of activated carbon and bentonite for adsorption of amoxicillin from wastewater: mechanisms, isotherms and kinetics. *Water Res.* 2015;43(9):2419-30.
- Ahmed MJ, Theydan SK. Microporous activated carbon from siris seed pods by microwave-induced KOH activation for metronidazole adsorption. *J Anal App Pyrol.* 2013;99: 101-9.
- Zhang D, Yin J, Zhao J, Zhu H, Wang C. Adsorption and removal of tetracycline from water by petroleum coke-derived highly porous activated carbon. *J Environ Chem Eng.* 2015;3(3):1504-12.
- Aksu Z, Tunç Ö. Application of biosorption for penicillin G removal: comparison with activated carbon. *Process Biochem.* 2005;40(2):831-47.
- Ahmed MJ, Theydan SK. Fluoroquinolones antibiotics adsorption onto microporous activated carbon from lignocellulosic biomass by microwave pyrolysis. *J Taiwan Inst Chem Eng.* 2015;45(1):219-26.
- Ahmed MB, Zhou JL, Ngo HH, Guo W. Adsorptive removal of antibiotics from water and wastewater: Progress and challenges. *Sci Total Environ.* 2015;532:112-26.
- Acosta R, Fierro V, Yuso AM, Nabarlaz D, Celzard A. Tetracycline adsorption onto activated carbons produced by KOH activation of tyre pyrolysis char. *Chemosphere.* 2016;149:168-76.
- Mansouri H, Carmona RJ, Gomis-Berenguer A, Souissi-Najar S, Ouederni A, Ania CO. Competitive adsorption of ibuprofen and amoxicillin mixtures from aqueous solution on activated carbons. *J Colloid Interf Sci.* 2015;449:252-60.
- Balarak D, Azarpira H, Mostafapour FK. Adsorption isotherm studies of tetracycline antibiotics from aqueous solutions by maize stalks as a cheap biosorbent.

- Int J Pharm Technol. 2016;8(3):16664-75.
30. Mahvi AH, Mostafapour FK. Biosorption of tetracycline from aqueous solution by azolla filiculoides: equilibrium kinetic and thermodynamics studies. *Fresenius Environ. Bull.* 2018;27(8):5759-67.
 31. Martins AC, Pezoti O, Cazetta AL, Bedin KC, Yamazaki DAS, Bandoch GFG, et al. Removal of tetracycline by NaOH-activated carbon produced from macadamia nut shells: kinetic and equilibrium studies. *Chem Eng J.* 2015;260:291-9.
 32. Mihciokur H, Oguz M. Removal of oxytetracycline and determining its biosorption properties on aerobic granular sludge. *Environ Toxicol Pharmacol.* 2016;46:174-82.
 33. Mohammadi AS, Sardar M. The removal of penicillin G from aqueous solutions using chestnut shell modified with H₂SO₄: isotherm and kinetic study. *Iranian J Health Environ.* 2012;6:497-508.
 34. Lin Y, Xu S, Jia L. Fast and highly efficient tetracyclines removal from environmental waters by graphene oxide functionalized magnetic particles. *Chem Eng J.* 2013;225:679-85.
 35. Liu W, Zhang J, Zhang C, Ren L. Sorption of norfloxacin by lotus stalk-based activated carbon and iron-doped activated alumina: Mechanisms, isotherms and kinetics. *Chem Eng J.* 2011;171(2):431-8.
 36. Balarak D, Azarpira H. Rice husk as a Biosorbent for Antibiotic Metronidazole Removal: Isotherm Studies and Model validation. *Int J ChemTech Res.* 2016;9(7):566-573.
 37. Azarpira H, Mahdavi Y, Khaleghi O. Thermodynamic Studies on the Removal of Metronidazole Antibiotic by Multi-Walled Carbon Nanotubes. *Pharm Lett.* 2016;8(11):107-13.
 38. Balarak D, Azarpira H, Mostafapour FK. Study of the Adsorption Mechanisms of Cephalalexin on to Azolla Filiculoides. *Pharm Chem.* 2016;8(10):114-21.
 39. Zhang Z, Li H, Liu H. Insight into the adsorption of tetracycline onto amino and amino-Fe³⁺ functionalized mesoporous silica: Effect of functionalized groups. *J Environ Sci.* 2018;65:171-8.
 40. Lv JM, Ma YL, Chang X, Fan SB. Removal and removing mechanism of tetracycline residue from aqueous solution by using Cu-13X. *Chem Eng J.* 2015;273:247-53.
 41. Xiong W, Zeng G, Yang Z, Zhou Y, Zhang C, Cheng M, et al. Adsorption of tetracycline antibiotics from aqueous solutions on nanocomposite multi-walled carbon nanotube functionalized MIL-53(Fe) as new adsorbent. *Sci Total Environ.* 2018;627:235-44.
 42. Cao J, Xiong Z, Lai B. Effect of initial pH on the tetracycline (TC) removal by zero-valent iron: Adsorption, oxidation and reduction. *Chem Eng J.* 2018;343:492-9.
 43. Marzbali MH, Esmaili M, Abolghasemi H, Marzbali MH. Tetracycline adsorption by H₂PO₄-activated carbon produced from apricot nut shells: A batch study. *Process Saf Environ Prot.* 2016;102:700-9.
 44. Ahmed MJ. Adsorption of quinolone, tetracycline, and penicillin antibiotics from aqueous solution using activated carbons: Review. *Environ Toxicol Pharmacol.* 2017;50:1-10.
 45. Vona A, Martino F, Garcia-Ivars J, Picó Y, Mendoza-Roca JA, Iborra-Clar MI. Comparison of different removal techniques for selected pharmaceuticals. *J Water Process Eng.* 2015;5:48-57.
 46. Ahmadi S, Banach A, Kord Mostafapour F. Study survey of cupric oxide nanoparticles in removal efficiency of ciprofloxacin antibiotic from aqueous solution: Adsorption isotherm study. *Desalin Water Treat.* 2017;89:297-303.
 47. Balarak D, Mostafapour F, Bazrafshan E, Saleh TA. Studies on the adsorption of amoxicillin on multi-wall carbon nanotubes. *Water Sci Technol.* 2017;75(7):1599-606.
 48. Yu B, Bai Y, Ming Z, Yang H, Chen L, Hu X, et al. Adsorption behaviors of tetracycline on magnetic graphene oxide sponge. *Mater Chem Phys.* 2017;198:283-90.
 49. Adriano W. Adsorption of amoxicillin on chitosan beads: Kinetics, equilibrium and validation of finite bath models. *Biochem Eng J.* 2005;27(2):132-7.
 50. Ali I, AL-Othman ZA, Alwarthan A. Synthesis of composite iron nano adsorbent and removal of ibuprofen drug residue from water. *J Mol Liq.* 2016;219:858-64.
 51. Carabineiro S. Comparison between activated carbon, carbon xerogel and carbon nanotubes for the adsorption of the antibiotic ciprofloxacin. *Catal today.* 2012;186(1):29-34.
 52. Ou J, Mei M, Xu X. Magnetic adsorbent constructed from the loading of amino functionalized Fe₃O₄ on coordination complex modified polyoxometalates nanoparticle and its tetracycline adsorption removal property study. *J Solid State Chem.* 2016;238:182-8.
 53. Zhang Y, Jiao Z, Hu Y, Lv S, Fan H, Zeng Y. Removal of tetracycline and oxytetracycline from water by magnetic Fe₃O₄@graphene. *Environ Sci Pollut Res.* 2017;24(3):2987-95.
 54. Harja M, Ciobanu G. Studies on adsorption of oxytetracycline from aqueous solutions onto hydroxyapatite. *Sci Total Environ.* 2018;628:36-43.
 55. Fazlirad H. Preparation of magnetic multiwalled carbon nanotubes for an efficient adsorption and spectrophotometric determination of amoxicillin. *J IndEng Chem.* 2015;21:889-2.
 56. Chen Y, Wang F, Duan L, Yang H, Gao J. Tetracycline adsorption onto rice husk ash, an agricultural waste: Its kinetic and thermodynamic studies. *J Mol Liq.* 2016;222:487-94.
 57. Jafari M, Aghamiri SF, Khaghanic G. Batch adsorption of cephalosporins antibiotics from aqueous solution by means of multi-walled carbon nanotubes. *World appl Sci J.* 2011;14(11):1642-50.
 58. Ji L. Adsorption of tetracycline on single-walled and multi-walled carbon nanotubes as affected by aqueous solution chemistry. *Environ Toxicol Chem.* 2011;29(12):2713-9.
 59. Kim H, Hwang YS, Sharma VK. Adsorption of antibiotics and iopromide onto single-walled and multiwalled carbon nanotubes. *Chem Eng Technol.* 2014;255:23-7.
 60. Kim SH, Shon HK, Ngo HH. Adsorption characteristics of antibiotics trimethoprim on powdered and granular activated carbon. *J IndEng Chem.* 2010;16(3):344-9.
 61. Zhang CL, Li BY, Wang Y. Adsorption of norfloxacin from aqueous solution onto modified coal fly ash. *Chem Eng Technol.* 2010;33(6):969-72.
 62. Zhang CL, Qiao GL, Zhao F, Wang Y. Thermodynamic and kinetic parameters of ciprofloxacin adsorption onto modified coal fly ash from aqueous solution. *J Mol Liq.* 2011;163(1):53-6.
 63. Yahiaoui I, Aissani-Benissad F, Fourcade F, Amrane A. Removal of tetracycline hydrochloride from water based on direct anodic oxidation (Pb/PbO₂ electrode) coupled to activated sludge culture. *Chem Eng J.* 2013;221:418-25.
 64. Balarak D, Mahdavi Y, Azadi NA, Sadeghi SH. Isotherms and thermodynamic study on the biosorption of amoxicillin using canola. *Int J Anal Pharm Biomed Sci.* 2016;5(3):8-14.
 65. Parolo ME, Savini MC, Vallés JM, Baschini MT, Avena MJ. Tetracycline adsorption on montmorillonite: pH and ionic strength effects. *Appl Clay Sci.* 2008;40(14):179-186.
 66. Yaghmaeian K, Moussavi G, Alahabadi A. Removal of amoxicillin from contaminated water using NH₄Cl-activated carbon: Continuous flow fixed-bed adsorption and catalytic ozonation regeneration. *Chem Eng J.* 2014;236:538-44.
 67. Ghauch A, Tuqan A, Assi HA. Elimination of amoxicillin and ampicillin by micro scale and nano scale iron particles. *Environ Pollut.* 2009;157(5):1626-35.
 68. Malakootian M, Mahdavi Y, Sadeghi SH, Amirmahani N. Removal of antibiotics from wastewater by azolla filiculoides: kinetic and equilibrium studies. *Int J Anal Pharm Biomed Sci.* 2015;4(7):105-13.
 69. Chen WR, Huang CH. Adsorption and transformation of tetracycline antibiotics with aluminum oxide. *Chemosphere.* 2010;79:779-85.
 70. Liu H, Yang Y, Kang J, Fan M, Qu J. Removal of tetracycline from water by Fe-Mn binary oxide. *J Environ Sci.* 2012;24(2):242-7.
 71. Zhao Y, Geng J, Wang X, Gu X, Gao S. Adsorption of tetracycline onto goethite in the presence of metal cations and humic substances. *J Colloid Interface Sci.* 2011;361(1):247-51.
 72. Prado N, Ochoa J, Amrane A. Biodegradation and biosorption of tetracycline and tylosin antibiotics in activated sludge system. *Process Biochem.* 2009;44(11):1302-6.
 73. Kang J, Liu H, Zheng Y, Qua J, Chen PJ. Systematic study of synergistic and antagonistic effects on adsorption of tetracycline and copper onto a chitosan. *J Colloid Interface Sci.* 2010;344(1):117-25.
 74. Shao L, Ren Z, Zhang G, Chen L. Facile synthesis, characterization of a MnFe₂O₄/activated carbon magnetic composite and its effectiveness in tetracycline removal. *Mater Chem Phys.* 2012;135(1):16-24.
 75. Brigante M, Schul PC. Removal of the antibiotic tetracycline by titania and titania-silica composed materials. *J Hazard Mater.* 2011;192(3):1597-608.
 76. Balarak D, Mostafapour FK, Akbari, H. Adsorption of Amoxicillin Antibiotic from Pharmaceutical Wastewater by Activated Carbon Prepared from Azolla filiculoides. *Br J Pharm Res.* 2017;18(3):1-10.
 77. Balarak D, Mostafapour FK, Joghtaei A. Thermodynamic Analysis for Adsorption of Amoxicillin onto Magnetic Carbon Nanotubes. *Br J Pharm Res.* 2017;16(6):1-10.
 78. Balarak D, Mostafapour FK, Joghataei A. Experimental and Kinetic Studies on Penicillin G Adsorption by Lemna minor. *Br J Pharm Res.* 2016;9(5):1-10.
 79. Zhao Y, Gu X, Gao S, Geng J, Wang X. Adsorption of tetracycline (TC) onto montmorillonite: cations and humic acid effects. *Geoderma.* 2012;183:12-8.
 80. Li Z, Schulz L, Ackley C, Fenske N. Adsorption of tetracycline on kaolinite with pH-dependent surface charges. *J Colloid Interface Sci.* 2010;351(1):254-60.
 81. Chang PH, Li Z, Jean J, Jiang W, Wang CJ, Lin KH. Adsorption of tetracycline on 2:1 layered non-swelling clay mineral illite. *Appl Clay Sci.* 2012;67:158-63.
 82. Chang PH, Li Z, Jiang WT, Jean JS. Adsorption and intercalation of tetracycline by swelling clay minerals. *Appl Clay Sci.* 2009;46(1):27-36.
 83. Godosa I, Munozb R, Guieyssea B. Tetracycline removal during wastewater treatment in high-rate algal ponds. *J Hazard Mater.* 2012;229:446-9.
 84. Liu P, Liu W, Jiang H, Chen J, Li W, Yu H. Modification of bio-char derived from fast pyrolysis of biomass and its application in removal of tetracycline from aqueous solution. *Bioresour Technol.* 2012;121:235-40.
 85. Homem V, Santos L. Degradation and removal methods of antibiotics from aqueous matrices- a review. *J Environ Manage.* 2011;92(10):2304-47.
 86. Balarak D, Mostafapour FK. Batch Equilibrium, Kinetics and Thermodynamics Study of Sulfamethoxazole Antibiotics Onto Azolla filiculoides as a Novel

- Biosorbent. *Br J Pharm Res.* 2016;13(2):1-10.
87. Balarak D, Mahdavi Y, Mostafapour FK. Application of Alumina-coated Carbon Nanotubes in Removal of Tetracycline from Aqueous Solution. *Br J Pharm Res.* 2016;12(1):1-11.
 88. Ji LG, Chen W, Duan L, Zhu D. Mechanisms for strong adsorption of tetracycline to carbon nanotubes: a comparative study using activated carbon and graphite as adsorbents. *Environ Sci Technol.* 2009;43(7):2322-7.
 89. Gao Y, Li Y, Zhang L, Huang H, Hu J, Shah SM, *et al.* Adsorption and removal of tetracycline antibiotics from aqueous solution by graphene oxide. *J Colloid Interface Sci.* 2012;368(1):540-6.
 90. Rahardjo A, Susanto M, Kurniawan A, Indraswati N. Modified Ponorogo bentonite for the removal of ampicillin from wastewater. *J Hazard Mater.* 2011;190(1-3):1001-8.
 91. Chang PH, Li Z, Yu TL, Munkhbayer S, Kuo TH, Hung YC. Sorptive removal of tetracycline from water by palygorskite. *J Hazard Mater.* 2009;165(1-3):148-55.
 92. Turku I, Sainio T, Patero E. Thermodynamics of tetracycline adsorption on silica. *Environ Chem Lett.* 2007;5(4):225-8.
 93. Balarak D, Mansouri AH, Chandrika K. Adsorption characteristics of metronidazole from industrial wastewater onto polyaniline nanocomposite. *Int J Life Sci Pharm Res.* 2019;9(4):49-58.
 94. Balarak D, Chandrika K. Investigating the Efficiency of Zirconium Oxide Nanoparticles in Removal of Ciprofloxacin from Aqueous Environments. *Int J Life Sci Pharm Res.* 2020;10(1):17-22.
 95. Faghihian H, Amini MK, Nezamzadeh AR. Cerium uptake by zeolite A synthesized from natural clinoptilolite. *J Rad Nucl Chem.* 2005;264(3):577-82.
 96. Ahmed MJ, Theydan SK. Microwave assisted preparation of microporous activated carbon from siris seed pods for adsorption of metronidazole antibiotic. *Chem Eng J.* 2013;214:310-8.
 97. Moussavi G, Alahabadi A, Yaghmaeian K, Eskandari M. Preparation, characterization and adsorption potential of the NH₄Cl-induced activated carbon for the removal of amoxicillin antibiotic from water. *Chem Eng J.* 2013;217:119-28.
 98. Ocampz PR, Leyva RR, Rivera UJ, Flores CJV, Sánchez PM. Modeling adsorption rate of tetracyclines on activated carbons from aqueous phase. *Chem Eng Res Design.* 2015;104:579-88.
 99. Mostafapour FK, Baniyasi M. Removal of Ciprofloxacin from of Pharmaceutical Wastewater by Adsorption on SiO₂Nanoparticle. *J Pharm Res Int.* 2018;25(6):1-9.
 100. Balarak D, Mostafapour FK, Joghataei A. Kinetics and mechanism of red mud in adsorption ciprofloxacin in aqueous solution. *Biosci Biotechnol Res commun.* 2017;10(1):241-8.
 101. Torres PJ, Gérente C, Andrés Y. Sustainable Activated carbons from agricultural residues dedicated to antibiotic removal by adsorption. *Chinese J Chem Eng.* 2012;20(3):524-9.
 102. Carabineiro SAC, Thavorn-Amornsri T, Pereira MFR. Adsorption of ciprofloxacin on surface-modified carbon materials. *Water Res.* 2012;45(15):4583-91.
 103. Xie HJ, Liu WF, Zhang J, Zhang CL, Ren L. Sorption of norfloxacin from aqueous solutions by activated carbon developed from *Trapanatans husk*. *Sci. China Chem.* 2011;54(5):835-43.
 104. Xu XR, Li XY. Sorption and desorption of antibiotic tetracycline on marine sediments. *Chemosphere.* 2010;78(4):430-6.
 105. Brinzila CI, Pacheco MJ, Ciriaco L, Ciobanu RC, Lopes A. Electro degradation of tetracycline on BDD anode. *Chem Eng J.* 2011;209:54-61.
 106. Yahya MA, Al-Qodah Z, Zanariah CW. Agricultural bio-waste materials as potential sustainable precursors used for activated carbon production: a review. *Renew Sustain Energy Rev.* 2015;46:218-35.
 107. Yang W, Lu Y, Zheng F, Xue X, Li N, Liu D. Adsorption behavior and mechanisms of norfloxacin onto porous resins and carbon nanotube. *Chem Eng J.* 2012;179:112-8.
 108. Li K, Ji F, Liu Y, Tong Z, Zhan X, Hu Z. Adsorption removal of tetracycline from aqueous solution by anaerobic granular sludge: equilibrium and kinetic studies. *Water Sci Technol.* 2013;67(7):1490-6.
 109. Liu H, Ning W, Cheng P, Zhang J, Wang Y, Zhang C. Evaluation of animal hairs-based activated carbon for sorption of norfloxacin and acetaminophen by comparing with cattail fiber-based activated carbon. *J Anal Appl Pyrol.* 2013;101:156-65.
 110. Li X, Chen S, Fan X, Quan X, Tan F, Zhang Y, *et al.* Adsorption of ciprofloxacin, bisphenol and 2-chlorophenol on electrospun carbon nanofibers: in comparison with powder activated carbon. *J Colloid Interf Sci.* 2015;447:120-7.
 111. Sun Y, Yue Q, Gao B, Wang B, Li Q, Huang L, *et al.* Comparison of activated carbons from *arundo donax* Linn with H₂P₂O₇ activation by conventional and microwave heating methods. *Chem Eng J.* 2012 192:308-14.
 112. Lian F, Song Z, Liu Z, Zhu L, Xing B. Mechanistic understanding of tetracycline sorption on waste tire powder and its chars affected by Cu²⁺ and Ph. *Environ Pollut.* 2013;178:264-70.
 113. Cong Q, Yuan X, Qu J. Removal of antibiotics by carbon nanotubes. *Water Sci Technol.* 2013;68(8):1679-87.
 114. Daghri R, Drogui P. Tetracycline antibiotics in the environment. *Environ Chem Lett.* 2013;11(3):209-27.
 115. Derakhshan Z. Removal methods of antibiotic compounds from aqueous environments - A review. *J Environ Health Sustainable Dev.* 2016;1(1):43-62.

Article History: Submission Date : 04-02-2020; Revised Date : 29-02-2020; Acceptance Date : 18-03-2020.

Cite this article: Balarak D, Khatibi AD, Chandrika K. Antibiotics Removal from Aqueous Solution and Pharmaceutical Wastewater by Adsorption Process: A Review. *Int. J. Pharm. Investigation.* 2020;10(2):106-11.